THE BRYOLOGIST

Quarterly Journal of the American Bryological and Lichenological Society

Editor
WILLIAM D. REESE

Volume 77
1974

Table of Contents

ANDERSON, R. A. Additions to the lichen flora of North America—III 41

——— & M.-B. CARMER. Rediscovery of *Lecidea populina* in Colorado 82

——— & ———. Additions to the lichen flora of Colorado 216

BAKER, R. G. Review 277

BOWERS, F. D. The mosses reported from Costa Rica 150

BOWLER, P. A. & P. W. RUNDEL. The *Ramalina intermedia* complex in North America 617

BRASSARD, G. R. Review 497

BURR, R. J., B. G. BUTTERFIELD & C. HÉBANT. A correlated scanning and transmission electron microscope study of the water-conducting elements in the gametophytes of *Haplomitrium gibbsiae* and *Hymenophyton flabellatum* 612

CHOPRA, R. N. & SHEEL. Some new observations on the gametangia in Splachnaceae 623

COOMBES, A. J. & N. W. LEPP. The effect of Cu and Zn on the growth of *Marchantia polymorpha* and *Funaria hygrometrica* 447

CROSBY, M. R. Recent literature on mosses—97–99 ..... 88, 262, 481

CRUM, H. Review 103

CULBERSON, W. L. Recent literature on lichens—87–90 ..... 94, 267, 488, 648

———. Review 101

DEY, J. P. *Parmelia commensurata*, a lichen new to North America 250

DISRUD, D. T. Two mosses new to North Dakota 84

DOUGLAS, G. W. Lichens of the North Cascades Range, Washington 582

FREY, W. Die Lebermoose des Irans und Afghanistans 48

FULFORD, M. Recent literature on hepatics—67–69 ..... 85, 258, 478

GLIME, J. M. & R. E. CARR. Temperature survival of *Fontinalis novae-angliae* Sull. 17

GROSSMAN, H. H. & C. J. HILLSON. The effect of chronic low-level gamma radiation on *Polytrichum commune*. I. Growth and development of gametophores from irradiated sporocytes and spores 142

GUERKE, W. R. A floristic study of the Hepaticae and Anthocerotae of the Florida Parishes, Louisiana 593

HALE, M. E., JR. & P. G. PATWARDHAN. Two new species of *Hypotrachyna* (Lichenes) from India 637

HANCOCK, J. A. & G. R. BRASSARD. Phenology, sporophyte production, and life history of *Buxbaumia aphylla* in Newfoundland, Canada 501

HATTORI, S. Review 657

HEGEWALD, E. Beitrag zur Moosflora Perus. I. *Bryomanginia saint-pierrei*—neu für Südamerika 641

HERMANN, F. J. & H. ROBINSON. Additions to the bryophyte flora of Bolivia 643

HICKS, M. L. Marginal gemmae in *Frullania eboracensis* 460

HOFFMAN, G. R. Growth stimulation of *Marchantia polymorpha* from ilmenite basalt and volcanic ash 632

HOLLENSEN, R. H. Review 497
IRELAND, R. R. *Neomacounia*, a new genus in the Neckeraceae (Musci) from Canada 453
JOHNSEN, A. B. *Tayloria splachnoides* in Arizona 636
KALAITZAKIS, J. A. & N. S. MARGARIS. Quantitative analysis of soluble sugars from two lichen species 465
KITAGAWA, N. & T. KODAMA. A remarkable new species of *Acromastigum* (Hepaticae) with septate rhizoids and filamentous gemmae 57
KROG, H. *Parmelia ultralucens*, a new lichen species in subgenus *Amphigymnia* 253
LAZARENKO, A. S. Some considerations on the nature and behavior of the relict moss, *Desmatodon randii* 474
LEWINSKY, J. Some new or noteworthy mosses from west Greenland 74
———. The genera *Leskeella* and *Pseudoleskeella* in Greenland 601
LONGTON, R. E. Microclimate and biomass in communities of the *Bryum* association on Ross Island, continental Antarctica 109
MAGILL, R. E. Notes on Texas mosses III 645
———, F. SEABURY & D. M. J. MUELLER. Evaluation of the critical point drying technique and its application in systematic studies of mosses 628
MANUEL, M. G. A revised classification of the Leucodontaceae and a revision of the subfamily Alsioideae 531
MARGARIS, N. S. Free amino acid pools in *Cladonia pyxidata* and *Peltigera* sp. 77
———. Free amino acid pools of certain mosses from Greece 246
——— & J. KALAITZAKIS. Soluble sugars from some Greek mosses 470
MARTON, K. & M. GALUN. A new species of *Heppia* from the Arava Valley, Israel 239
MELSTROM, C. E., N. C. MARAVOLO & J. R. STROEMER. Endogenous gibberellins in *Marchantia polymorpha* and their possible physiological role in thallus elongation and orthogeotropic growth 33
MEYER, M. W., J. GREENBERG & S. TEDESCHI. Enzymes of the moss *Funaria hygrometrica* II: The isoenzymes of malate dehydrogenase 577
MILLER, N. G. Review 100
MOLENAAR, J. G. DE. *Dicranella riparia* new to Greenland 236
NASH, T. H., III. New and additional *Xanthoparmelia* from New Mexico 72
———. Two new species of *Xanthoparmelia* with diffractaic acid 234
———. Review 661
——— & W. A. WEBER. Additions to the lichen flora of Arizona II 472
NEWBERRY, G. The influence of a sulfate-process paper mill on corticolous lichens 561
OSORIO, H. S. *Placomaronea candelarioides*: Ampliación de su área de distribución geográfica 463
REDFEARN, P. L., JR. Reviews 279, 496
REESE, W. D. Reviews 103, 495
———. *Syrrhopodon quintasii* in the Americas 242
———. *Bryoxiphium norvegicum* in Arizona 252
———. Notes on Louisiana mosses—VI 467
REYES, C. R. Nota adicional acerca del catalogo de los liquenes de Venezuela 248

——— & F. Bowers. Contribución á la brioflora del Macarao, Venezuela I 241
——— & A. Skorepa. Contribución á la flora liquenologica del Macarao, Venezuela I 257
Rundel, P. W. Water relations and morphological variation in *Ramalina menziesii* Tayl. 23
———. Review 660
——— & P. A. Bowler. The lichen genus *Trichoramalina* 188
Schuster, R. M. Review 659
Selkirk, D. R. A fossil epiphyllous moss from the Australian Miocene 249
Seppelt, R. D. *Riccia crystallina* in south Australia 224
Sheard, J. W. The genus *Dimelaena* in North America north of Mexico 128
——— & M. E. Jonescu. A multivariate analysis of the distribution of lichens on *Populus tremuloides* in west-central Canada 514
Sigal, L. L. & D. R. Toren. New distribution records of lichens in California 469
Smiley, D. & C. J. George. Photographic documentation of lichen decline in the Shawangunk Mountains of New York 179
Snider, J. A. & F. D. Bowers. *Archidium*, a moss genus new to Mexico 639
Solberg, Y. J. Studies on the chemistry of lichens. XII. Chemical investigation of the lichen species *Xanthoria parietina* (L.) Th. Fr.—2 203
Steere, W. C. The occurrence of *Mnium spinosum* in North America. II 79
———. Review 103
———. *Grimmia* (*Coscinodon*) *arctolimnia*, a new species from Great Bear Lake, Northwest Territories, Canada 230
——— & G. R. Brassard. The systematic position and geographical distribution of *Fissidens arcticus* 195
——— & H. Inoue. *Fossombronia alaskana*, a new hepatic from arctic Alaska 63
——— & B. M. Murray. The geographical distribution of *Bryum wrightii* in arctic and boreal North America 172
Stringer, P. W. & M. H. L. Stringer. Studies on the bryophytes of southern Manitoba. VI. An ecological study of the bryophytes of coniferous forests in Bird's Hill Provincial Park 1
——— & ———. Seventeen lichens new to Manitoba 243
——— & ———. Air pollution and the distribution of epiphytic lichens and bryophytes in Winnipeg, Manitoba 405
——— & ———. A quantitative study of corticolous bryophytes in the vicinity of Winnipeg, Manitoba 551
Taylor, J., P. J. Hollingsworth & W. C. Bigelow. Scanning electron microscopy of liverwort spores and elaters 281
Traverse, A. Review 495
Weber, W. A. Review 276
Welch, W. H. The Hookeriaceae of Central America 328
Wetmore, C. M. Review 104
———. New type of soredium in the lichen family Heppiaceae 208
Whittier, H. O. & B. A. Whittier. List of mosses of southeastern Polynesia 427

Index to Volume 77 of The Bryologist[1, 2]

Acanthocladium costaricensis, 167; subnitidum, 167
Acarospora, 128-, 469-; chlorophana, 584; fuscata, 584; sinopica, 216; washingtonensis, 584
Acaulon, 63-; muticum var. rufescens, 645
Acidodontium floresianus, 157; megalocarpon, 157
Acrobolbus cinerascens, SEM of spores and elaters, 299, 304*
Acrocladium, 441
Acrocryphaea, generic characters, 533*
Acrocryphaea gardneri, 160; julacea var. costaricensis, 160
Acromastigum sect. Acromastigum, 57; sect. Exilia, 57; sect. Inaequilatera, 57; sect. Subcomplicata, 57; subg. Triandrophyllopsis, 57
Acromastigum **gemmiferum** sp. nov., 57, 58*, 60*; integrifolium, 57; obliquatum, 61; stenophyllum, 57
Acroporium brevicuspidatum, 441; lepinei, 441; pungens, 167; sigmatodontium, 441
Actinodontium portoricense, 332; sprucei, 163, 331; standleyi, 331, 332*
Adaptations of Ramalina menziesii, 29
Adelanthus pittierii, 644
Adelothecium bogotense, 163, 332
Aerobryopsis longissima, 437
Aerobryum vitianum, 437
Afghanistan, die Lebermoose, 48
Agyrophora lyngei, 584
Air pollution effects: on bryophyte distribution, 405; on corticolous lichens, 561, 564*, 565*; on lichen distribution, 405
Alectoria americana, 212; fremontii, 584; glabra, 584; jubata, 584; minuscula, 469, 584; nigricans, 584; ochroleuca, 584; oregona, 584; pubescens, 584; sarmentosa, 584
Alliance, Calamagrostion neglectae, 237
Aloina rigida, 467
Aloinella hamulus, 467
Alsia, generic characters, 533*
Alsia abietina, *545*; californica, 453, 539, 540*, 541*, distribution of, 534*; californica var. flagellifera, 539; circinalis, *546*; circinata, *545*; circinnata, *545*; longipes, 537; macounii, *545*
Amblystegium byssoides, 440; compactum, 475; juratzkanum, 8, 415, 556; juratzkanum union, 8; montanae, *602*; serpens, 7, 415, 556; varium, 415, 556, 644; varium var. varium, 6
Amino acids: in Cladonia pyxidata, 77; in Peltigera, 77
Amphidium, 230-
Amphiloma, *130*
Amphomorphism, 476
Anaptychia speciosa, 414; tropica, 472
Anastrophyllum minutum, 74-
Anderson, R. A.: Additions to the lichen flora of North America III, 41; and M.-B. Carmer: Rediscovery of Lecidea populina in Colorado, 82; and M.-B. Carmer: Additions to the lichen flora of Colorado, 216
Andreaea rupestris, 74-, 151; turgescens, 151
Aneura pinguis, 596
Anisothecium rufescens, 152; vaginatum, 152
Anoectangium aestivum, 154; condensatum, 154; euchloron, 154, 433; tapes, 433
Anomobryum costaricense, 157; filiforme, 157; plicatum, 157; semiovatum, 157
Anomodon minor, 415, 556, 645; nervosus, *602*; tectorum, *606*
Anoplolejeunea conferta, 644
Antarctica, map, 111*
Antarctica, microclimate and biomass of Bryum associations, 109
Anthelia, 63-; juratzkana, 236-
Antheridia, in Splachnaceae, 623
Anthoceros punctatus, 596
Antitrichia, generic characters, 533*
Antitrichia californica, 537; californica var. flagellifera, 539; curtipendula, 543; pseudocalifornica, 539; tenella, 543; tenella var. flagellarifera, 539
Aongstroemia jamaicensis, 152; julacea, 152
Apogamy, Desmatodon capsules, 475
Apospory, Desmatodon protonema, 475
Apothecia, methods for morphologic study, 129
Apotreubia nana, gametangia, 627

[1] Compiled by R. E. Stotler, Botany Department, Southern Illinois University, Carbondale, Illinois 62901.

[2] New scientific names are printed in **boldface** type; illustrations are indicated by an asterisk; synonyms, by italicized page numbers; minor entries, by a dash following the initial article page.

Aptychella americana, 167
Arachniopsis monocera, 57-
Archegonia, in Splachnaceae, 623
Archidium alternifolium, 639; amplexicaule, 639; cubense, 639; donnellii, 639; hallii, 639, 640*; hallii var. minus, 639; julaceum, 639; ohioense, 639; tenerrimum, 639; sect. Protobium, 639
Arctic, distribution of Bryum wrightii, 172
Arctic Alaska, a new hepatic, 63
Argentina, Placomaronea candelarioides in, 463
Arizona, lichen additions, 472; Bryoxiphium norvegicum in, 252; Tayloria splachnoides in, 636
Arthonia, 220, 414; caesia, 568; cyanea, 248; lapidicola, 216; mira, 248; radiata, 584; trilocularis, 248
Arthopyrenia analeptella, 584
Arthothelium, 41-, 220
Arthrocormus dentatum, 431; nadeaudii, 431; schimperi, 431
Asexual propagules of Buxbaumia aphylla, 510*
Aspicilia cinereorufescens, 216
Association table of Manitoba bryophytes, 7*
Associations: Calamagrosto-Ditrichetum, 237
Asterella bolanderi, 281-; californica, 281-; gracilis, 50; palmeri, 281-; persica, 49; tenella, 597
Asterella tenera, SEM of spores and elaters, 314, 316*, 318*
Asterothyrium lecophthalmum, 248; pittieri, 248
Athalamia hyalina, 50
Atractylocarpus costaricensis, 153; longisetus, 153
Atranorin, 41, 219, 250, 253, 619, 637
Atrichum androgynum, 169; hirtella, 169; oerstedianum, 169; polycarpum var. oerstedianum, 169; undulatiforme, 169; undulatum, 646
Aulacomnium, 623-; acuminatum, 195-; palustre, 8
Aulacomnium palustre-Sphagnum capillaceum union, 8
Australia, epiphyllous moss from the Miocene, 249; Riccia crystallina in, 224

Bacidia, 82-; akompsa, 516; albescens, 584; brasiliensis, 248; chlorococca, 566; declinis, 516; dimidiata, 248; fuscatula, 248; inundata, 216; populorum, 244, 414, 516; psychotriae, 248; stanhopeae, 248; vermifera, 516
Baeomyces absolutus, 257; rufus, 469, 584
Baker, R. G. (review), H. J. B. Birks, Past and Present Vegetation of the Isle of Skye. A Palaeoecological Study, 277
Barbella tenuissima, 161
Barbilophozia lycopodioides, 636-
Barbula, 6; bescherellei var. stenocarpa, 155; costaricensis, 155; cruegeri, 155; imperfecta, 644; pendula, 593-; rectifolia, 155; subulifolia, 155
Bartramia angustifolia, 158; costa-ricensis, 158; defoliata, 158; microstoma, 158; potosica, 158; sphaericarpa, 158; strumosa, 158
Bazzania trilobata, 597
Beltraminia, *130*
Berengeria sect. Placothallia, *130*
Bestia, generic characters, 533*
Biatorella kulshanensis, 584; microhaema, 244, 414; moriformis, 472; ochrophora, 82-
Bigelow, W. C. *See* Taylor, J., P. J. Hollingsworth, and W. C. Bigelow
Biomass estimation in mosses, 122
Blasia pusilla, 281-
Blasia pusilla, SEM of spores and elaters, 309*, 310
Blepharostoma trichophyllum, 6, 636-
Bolivia, *see* Latin America
Boreal, distribution of Bryum wrightii, 172
Bowers, F. D.: The Mosses Reported from Costa Rica, 150; *see* Reyes, C. R. and F. Bowers; *see* Snider, J. A. and F. D. Bowers
Bowler, P. A. and P. W. Rundel: The Ramalina intermedia complex in North America, 617; *see* Rundel, P. W. and P. A. Bowler
Brachymenium barbae-montis, 156; brachypelma, 156; bulbiferum, 156; capillare, 156; filiscens, 156; indicum, 433; jamesonii, 156; melanothecium, 433; pittieri, 156; spathulifolium, 156; speciosum, 156; standleyi, 157; svihlae, 157; systylium, 157; viviparum, 156
Brachythecium acuminatum, 6, 556; albofla-vens, 165; campestre, 7, 415, 556; collinum, 6, 415, 555; costaricense, 165; curtum, 6; delicatum, 165; flexiventrosum, 165; laxireticulatum, 165; longipes, 440; rutabulum, 8, 415, 556; salebrosum, 8, 413, 556; stereopoma, 165; tearapense, 440; velutinum, 415; velutinum var. velutinum, 6
Brassard, G. R. (review), G. Philippi, Moosflora und Moosvegetation des Freeman-Sund-Gebietes (Südost-Spitzbergen), 497; *see* Hancock, J. A. and G. R. Brassard; *see* Steere, W. C. and G. R. Brassard
Breutelia brittoniae, 158; chrysea, 158; deflexifolia, 158; eugeniae, 435; jamaicensis, 158; subarcuata, 158; tomentosa, 158
Brotherella minutula, 168
Bryoandersonia illecebra, 646

Bryoerythrophyllum campylocarpum, 155; recurvirostrum, 6, 155
Bryomanginia saint-pierrei, 641, 642*
Bryosedgwickia densa, 166; subdensa, 166
Bryoxiphium norvegicum subsp. norvegicum var. norvegicum, 252
Bryum associations, relation between microclimate and physiology, 119
Bryum, 415, 556, 623-; andicola, 157; angustirete, 6; antarcticum, 17-, 110; argenteum, 74-, 110, 157; argenteum var. costaricense, 157; argenteum var. lanatum, 157, 434; australe, 433; bigibbosum, 434; billardieri, 434; caespiticium, 6; capillare, 157; chryseum, 157, 644; coronatum, 157, 644; creberrimum, 6; cruegeri, 157; erythroneuron, 157; globosum, 173; globosum var. ruberrimum, 173; insolitum var. brachycarpum, 157; julaceus, 157; larminati, 434; leptothecium, 434; limbatum, 157; longii, 173; megalostegium, 434; mnioides, 157; oerstedianum, 156; pallens, 238; pallescens, 6; procerum, 157; pseudotriquetrum var. pseudotriquetrum, 6; rapaense, 434; rosulicoma, 157; rubrifolium, 157; ruderale, 468; rufo-limbatum, 157; semiovatum, 157; soboliferum, 644; sordidum, 157; spectabile, 156; stenotrichum var. stenotrichum, 6; truncorum, 434; weberaceum, 434; wrightii, 63-; wrightii, distribution in North America, 172, 174*, 176*
Buellia, 82-, 128-; aethalea, 217; alboatra, 584; badia, 472; disciformis, 584; epigaea, 584; fimbriata, 136; jugorum, 41, 217; lacteoidea, 217; papillata, 584; punctata var. polyspora, 567; radiata, *135*; semitensis, 41, 217; spuria, 217; stellulata, 128-; triphragmioides, 42, 217; wahlenbergii, 217; zahlbrucknerii, 584
Burr, R. J., B. G. Butterfield, and C. Hébant: A correlated scanning and transmission electron microscope study of the water-conducting elements in the gametophytes of Haplomitrium gibbsiae and Hymenophyton flabellatum, 612
Butterfield, B. G. *See* Burr, R. J., B. G. Butterfield, and C. Hébant
Buxbaumia aphylla, asexual propagules, 510*; life cycle, 501; phenology, 501, 512*; sporophyte development, 501, 502*, 503*
Byssolecania fumosonigricans, 248
Byssoloma leucoblepharum, 248; rotuliforme, 248; tricholomum, 248

Calicium trabinellum, 472; viride, 584
California, lichen additions, 469
Callicladium haldanianum, 6
Callicostella aquatica, 333, 334*; bernoullii, 335; chlorina, 438; ciliata, 336; depressa, 336; grossiretis, 334*, 336; hondurensis, 334*, 337; nukahivensis, 438; oblongifolia, 438; oerstediana, 163, 338*, 339; oerstedii, 339; pallida, 163, 339; papillata, 438; rivularis, 340; scabriseta, 340; tenerrima, 163, 338*, 341; vatteri, 338*, 341; vesiculata, 438
Calliergon giganteum, 12; giganteum var. giganteum, 7; stramineum, 236-
Calobryum rotundifolium, 281-
Calomnion denticulatum, 434; nadeaudii, 434; schistostegiellum, 434
Caloplaca, 82-; atrosanguinea, 584; aurantiaca, 516; bryochrysion, 42; cerina, 414, 516, 584; cinnamomea, 42; cirrochroa, 217; citrina, 244, 414; darbishirei, 109-; decipiens, 472; epithallina, 472; ferruginea, 585; herbidella, 42; holocarpa, 244, 414, 569, 585; jungermanniae, 473, 585; murorum, 585; pyracea, 516; squamosa, 473; stillicidiorum, 585; ulmorum, 244, 414, 516, 567
Calymperes, 242-; aduncifolium, 432; aongstroemii, 432; donnellii, 154; lonchophyllum, 154; mooreae, *433*; moorei, 432; nicaraguense, 154; nukahivense, 432; porrectum, 432; pseudopodianum, 432; quaylei, 432; richardii, 154; tahitense, 432; tenerum, 432; tuamotuense, 432; vriesei, 432
Calypogeia fissa, 51, 596; sullivantii, 596
Calyptothecium crispulum, 437; praelongum, 437; turgescens, 162; urvilleanum, 437
Camptothecium sp., sugars, 471*
Campylium chrysophyllum, 7; chrysophyllum var. chrysophyllum, 646; hispidulum, 8, 415, 555, 647; hispidulum var. sommerfeltii, 165; stellatum, 8, 74-
Campylodontium drepanioides, 166; onustum, 166
Campylopodium euphorocladum, 430; pusillum, 152
Campylopus aoraiensis, 430; arctocarpus, 152; atratus, *152*; blumii, 430; chrismarii, 152; concolor, 152; costaricensis, 152; donnellii, 152; dozyanus, 430; falcatulus, 152; filifolius, 152; flexuosus, 152; fragilis, 152; guatemalensis, 152; harrisii, 152; harrisii var. longifolius, 152; hellerianus, 152; hoffmannii, 152; introflexus, 152, 430; jamesonii, 152; nadeaudianus, 430; nudicaulis, 430; obscurus, 430; oerstedianus, 152; **paramoensis** nom. nov., 152; poasensis, 152; porphyreodictyon, 153; richardii, 153, 430; savannarum, 153; schimperi, 77; standleyi, 152; standleyi var. lutescens, 152; straminifolius, 152; subconcolor, 153; subfulvus, 644; subleucogaster, 152; subprolif-

erus, 152; sulfureus, 430; trichophorus, 153; umbellatus, 430; vernieri, 430
Canada: bryophytes, lichens, and air pollution, 405; lichens new to Manitoba, 243; **Neomacounia** gen. nov., 453; new species of Grimmia from the Northwest territories, 230; phenology of Buxbaumia aphylla, 501; west central, distribution of lichens on Populus tremuloides, 514
Candelaria concolor, 244, 414, 567; concolor var. effusa, 567
Candelariella aurella, 516, 585; cf. terrigena, 585; vitellina, 414; xanthostigma, 217
Carmer, M.-B. *See* Anderson, R. A. and M.-B. Carmer
Carr, R. E. *See* Glime, J. M. and R. E. Carr
Catagonium gracile, 441; politum, 162
Catharinaea hirtella, 169; oerstedianum, 169; polycarpa, 169
Catillaria atropurpurea, 244, 414; bouteillei, 585; glauconigricans, 516
Central America, *see* Latin America
Cephalozia ambigua, 236-; bicuspidata, 238; catenulata, 6, 596; connivens, 281-, 596; lunulifolia, 51, 596; media, 6
Cephaloziella arctica, 236-; byssacea, 632-; hyalina, 596; obliqua, 596; rhizantha, 596; rubella, 6; rubella subsp. arctogena, 74-
Ceratodon purpureus, 8, 74-, 109-, 172-, 415, 557; stenocarpus, 151
Ceratolejeunea laetefusca, 597
Cetraria chlorophylla, 585; ciliaris, 585; commixta, 585; cucullata, 585; ericetorum, 585; hepatizon, 585; idahoensis, 585; islandica, 585; juniperina, 585; nivalis, 585; pinastri, 585; platyphylla, 585; subalpina, 585
Cetrelia cetrarioides, 585; olivetorum, 212
Chaenotheca ferruginea, 218
Chaetephora pendula, *402*
Chaetomitrium tahitense, 438
Chandonanthus squarrosus, SEM of spores and elaters, 295, 298*
Cheilolejeunea clausa, 597; rigidula, 597
Chemistry of lichens, 203
Chiloscyphus pallescens, 7, 597; polyanthos, 51, 281-; polyanthos var. rivularis, 281-
Chiodecton sanguineum, 257
Chopra, R. N. and Sheel: Some new observations on the gametangia in Splachnaceae, 623
Chorisodontium setaceum, 153; speciosum, 153
Chromosome numbers, Desmatodon randii, 476
Cinclidium arcticum, 195-; stygium, 74-
Cladina arbuscula, 585; mitis, 585; rangiferina, 585
Cladonia pyxidata, sugars, 465*, 466*, 467*
Cladonia, 561-; aggregata, 257; bacillaris, 585; bellidiflora, 585; capitata, 585; cariosa, 585; carneola, 585; cenotea, 212; chlorophaea, 569, 586; coccifera, 586; coniocraea, 586; cornuta, 586, deformis, 473, 586; ecmocyna, 586; fimbriata, 586; floerkeana, 586; furcata, 586; gonecha, 586; gracilis, 586; hookeri, 586; lepidota, 586; macilenta, 586; multiformis, 586; phyllophora, 586; pityrea, 586; pyxidata, 586; squamosa, 569, 586; subsquamosa, 586; subulata, 586; verticillata, 586
Clasmatodon parvulus, 646
Clevea hyalina, 281-
Climacium dendroides, 8
Climate, of Trichoramalina habitats, 190*
Cluster analysis of bryophyte stands, 4, 6*
Cnestrum, 74-; alpestris, 77
Coccocarpia cronia, 257; epiphylla, 248
Coleochaetium standleyi, 158
Collema, 208-; subfurvum, 212; texanum, 473
Cololejeunea cardiocarpa, 597; contractiloba, 597; minutissima subsp. minutissima, 597; minutissima subsp. myriocarpa, 597; rossettiana, 52; setiloba, 597
Colorado, additions to the lichen flora, 216; Lecidea populina in, 82
Coniferous forests, an ecological study of the bryophytes of, 1
Coniocybe furfuracea, 218
Conocephalum conicum, 50, 281-, 596
Conostomum pentastichum, 158; tetragonum, 74-
Constictic acid, 639
Coombes, A. J. and N. W. Lepp: The effect of Cu and Zn on the growth of Marchantia polymorpha and Funaria hygrometrica, 447
Cornicularia aculeata, 586; normoerica, 469, 586
Corsinia coriandrina, 281-
Corticolous lichens of Winnipeg, 414*
Coscinodon trinervis, 233
Costa Rica, *see* Latin America
Critical point drying technique in mosses, 628
Crosby, M. R.: Recent literature on mosses, 97, 88; 98, 262; 99, 481
Crossomitrium acuminatum, 342, 343*; herminieri, 344; heterodontium, 164; oerstedianum, 164, 343*, 344; orbiculatum, 345; patrisiae, 164, 345; scabrisetum, 343*, 345; wallisii, 164, 346
Crum, H. (review) O. Degener, I. Degener, and H. Hörmann: Mosses of Hawaii, 103
Cryphaeaceae, 532
Cryphaea, generic characters, 533*
Cryphaea californica, *539*; filiformis, 160;

glomerata, 646; helictophylla, 437; nigrescens, 437; nitidula, 160; patens, 160; tahitica, 436; tenella, 436
Cryptochila grandiflora, 644
Cryptodontopsis, generic characters, 533*
Ctenidiadelphus cylindricarpus, 168
Ctenidium malacodes, 168; molluscum, 647; stellulatum, 442
CULBERSON, W. L.: Recent literature on lichens, 87, 94; 88, 267; 89, 488; 90, 648; (review) B. W. Ferry, M. S. Baddeley, and D. L. Hawksworth (editors) Air Pollution and Lichens, 101
Culture: Funaria hygrometrica protonemata, 577; Marchantia polymorpha, 34
Cuspidatula monodon, SEM of spores and elaters, 295, 300*
Cyathophorella tahitensis, 439
Cyathophorum bulbosum var. tahitense, 439; tahitense, 439
Cyclodictyon albicans, 163, 346; blumeanum, 439; brittonae, 163, 348*, 349; cuspidatum, 163, 349; erubescens, 350; humectatum, 350; jamesii, 163, 348*, 350; maxonii, 348*, 350; roridum, 351; rubrisetum, 163, 351; varians, 351
Cylindrothecium, 440; turgidum, *440*
Cynodontium strumiferum, 74-
Cyphelium inquinans, 586; lucidum, 218
Cyptodon, generic characters, 533*

Dactylina arctica, 586
Daltonia aristifolia, 163; gracilis, 163, 352; lindigiana, 163, 353, 354*; lindigii, 353; longifolia, 163, 353; pulvinata, 163, 353, 354*; sphaerica, 439; stenophylla, 163, 355; tenuifolia, 163, 354*, 356
Dasymitrium nadeaudii, *435*
Dendralsia, *545*
Dendroalsia, generic characters, 533*
Dendroalsia abietina, 541*, 544*, 545; abietina, distribution, 534*; circinalis, *546*; circinnata, *546*; donna, *546*
Dendrocryphaea, generic characters, 533*
Dendropogonella, generic characters, 533*
Dendropogonella, 437; rufescens, 160
Dermatocarpon acarosporoides, 473; fluviatile, 586; miniatum, 586; waltheri, 43
Desmatodon cernuus, 474; convolutus, 63-; heimii, 172-, 475; leucostoma, 172-; oxneri, 474; randii, 474; ucrainicus, 475
Desmatodon convolutus, SEM and use of critical point drying technique, 630*
Desmazieria, 189; melanothrix, *194*
DEY, J. P.: Parmelia commensurata, a lichen new to North America, 250
Diaspore types, 208
Dicnemon banksii, 431; rugosum, 431
Dicranella harrisii, 152; herminieri, 152; heteromalla, 152; heteromalla var. pittieri, 152; hilariana, 152; hilariana var. leptorhyncha, 152; hillebrandii, 430; hochreutineri, 430; leptorhyncha, 152; perrottetii, 152; riparia, 236-; riparia, distribution, 236*, 237*; rufescens, 152; rufiseta, 431; subinclinata, 152; standleyi, 152; tonduzii, 152; vaginata, 152
Dicranodontium meridionale, 153
Dicranoloma braunii, 431; brittonae, 153; rufifolium, 431; setaceum, 153
Dicranum costaricense, 153; frigidum, 153; oerstedianum, 152; pittieri, 153; polysetum, 6; strigulosum, 153
Didymodon campylocarpus, 155; fusco-viridis, 155
Diffractaic acid, 234, 251
Dimelaena, of North America, 128
Dimelaena angelica, 129; californica, 131; californica, chemical variants, 132*; californica, distribution map, 133*; chrysomelaena, 129; diffractella, 137*, 139; mulleri, 132; oreina, 132, 137*; oreina, characters, 135*; oreina, fumarprotocetraric acid variant, distribution, 133*; oreina, stictic-norstictic acid variant, distribution, 133*; radiata, 135, 137*; radiata, distribution, 139*; thysanota, 137*, 138; thysanota, distribution, 139*
Dimerella dilucida, 248; diluta, 218; epiphylla, 248
Diplophyllum apiculatum, 281-
Diploschistes actinostomus, 586; actinostomus var. brunneus, 473; scruposus, 586
Dirinaria picta, 257
DISRUD, D. T.: Two mosses new to North Dakota, 84
Distichium capillaceum, 6
Distichophyllum cuspidatum, 439; nadeaudii, 439; spathulatum, 439; tahitense, 439
Distribution of: Archidium hallii, 640*; Bryomanginia saint-pierrei, 642*; Bryum wrightii in North America, 176*; corticolous lichens and mosses of Winnipeg, 416*, 418*; Dendroalsia and Alsia, 534*; Dicranella riparia, 236*; Dimelaena in North America, 133*, 139*; Fissidens arcticus, 199*; Fossombronia in Alaska, 70*; Isopterygium muellerianum, 76*; Leskeella nervosa, 603*; Minum spinosum, 80*; Parmelia ultralucens, 255*; Placomaronea candelarioides en América del Sur, 464*; Pseudoleskeella catenulata, 605*; Pseudoleskeella papillosa, 610*; Pseudoleskeella tectorum, 608*; Ramalina intermedia in North America, 618*; Ramalina menziesii in California, 24*; Ramalina petrina, 622*; Rhabdoweisia crispata, 76*; Riccia crystal-

lina in south Australia, 225*; Trichoramalina, 193*
Ditrichum crinale, 644; pallidum, 646; pusillum, 646; cf. pusillum, 236-; steyermarkii, 151
Divaricatic acid, 619
Douglas, G. W.: Lichens of the North Cascades Range, Washington, 582
Dozya, generic characters, 533*
Drepanocladus aduncus var. aduncus, 6; aduncus var. kneiffii, 6; aduncus var. polycarpus, 8, 555; exannulatus, 236-; fluitans var. fluitans, 6; revolvens, 74-; uncinatus, 236-, 555; uncinatus var. uncinatus, 6
Dumortiera hirsuta, 597

Echinoplaca strigulacea, 248
Ecological studies: air pollution calculations of SO_2 dispersal patterns, 562*; air pollution effects on bryophytes and lichens, 405; air pollution effects on lichens, 561; air pollution indices, 413; burning and the distribution of Buxbaumia aphylla, 502; copper and bryophyte growth, 447; correlation of bryophyte stands with vascular strata, 11; correlation of bryophyte stands with soil, 11, 13*; cover of corticolous bryophytes, 554*; cover, recording for epiphytes, 412; microclimate and biomass of Bryum associations, 109; microclimate determination, 111; moisture indices of bryophyte unions, 9*; multivariate analysis of lichen distribution, 514; *p*H measurements on bark, 563; phenology, 503, 505*; precipitation and spore diameter in Riccia crystallina, 228*; quantitative study of corticolous bryophytes, 551, 557*; sampling of corticolous bryophytes, 3, 553; similarity indices in lichen populations, 570*, 571*; water content, measurement in lichens, 25; zinc and bryophyte growth, 447
Ecology of: bryophytes of coniferous forests, 1; Dicranella riparia, 237; Fossombronia alaskana, 69; Hepaticae in Florida Parishes, Louisiana, 593; Manitoba bryophytes, dendrogram, 6*; Ramalina menziesii, 23; Trichoramalina, 190*
Ectropothecium amabile var. brevifolium, 168; apiculatum, 168; bryifolium, 442; costaricense, 168; eccremocladum, 442; fosbergii, 442; polyandrum, 442; sandwichense, 442; sodale, 442; subdenticulata, 168; subdenticulata var. latifolium, 168; venustulum, 442
Elaters, scanning electron microscopy of, *see* SEM studies of elaters
Encalypta procera, 6; rhabdocarpa, 601-
Entodon aurescens, 166; bernoullii, 166; beyrichii, 166; erythropus, 166; jamesonii, 166; pallescens, 166; pallidus, 440; polycarpus, 166; seductrix, 646; solanderi, 440; solanderi var. **turgidus** stat. nov., 440; stenocarpus, 166; turgidus, *440*
Entosthodon bonplandii, 156
Enzymes: acid phosphatase tests, 615; extractions in moss protonemata, 577; in Funaria hygrometrica, 577, 578*, 579*
Ephemeropsis, 249*; tjibodensis, 250; trentepohlioides, 250
Ephemerum, 195-
Epipterygium, 439, 442; immarginatum, 156
Eriodon radicalis, 166
Eriopus pacificus, 439
Erythrodontium densum, 166; longisetum, 166, 241; squarrosum, 166; subdensum, 166
Eucatagonium politum, 162
Eurhynchium exasperatum, 166; hians, 646; pulchellum, 8, 415, 555; vagans, 440
Eustichia spruceana, 157
Evernia mesomorpha, 567; prunastri, 587
Evernic acid, 619
Exodictyon dentatum, 431; nadeaudii, 431
Extractions of: endogenous gibberellins, 34; free amino acids, 77; lichen substances, 203

Fabronia flavinervis, 164; polycarpa, 164
Felipponea, generic characters, 533*
Fissidens sect. bryoidium, 195-; sect. semilimbidium, 195-
Fissidens aoraiensis, 429; arcticus, 63-, 195, 198*, 199*; asplenioides, 151; barbaemontis, 151; bryoides, 196; bryoides subsp. longifolius, 151; bryoides subsp. viridulus, 151; clarkii, 429; costaricensis, 151; diplodus, 151; dixonianus, 429; elegans, 151; exiguus, 197; fautauae, 429; flexinervis, 151; garberi, 151; impar, 197; jaiorum, 429; kegelianus, 151; longifolius, 151; mangarevensis, 429; mollis, 151; moorea, 430; nadeaudii, 430; nanobryoides, 430; oerstedianus, 151; osmundoides, 6; pellucidus, 151; philonotulus, 430; polypodioides, 151, 647; prionodes, 151; raiatensis, 430; reclinatulus, 151; repandus, 151; steyermarkii, 151; taiarapuensis, 430; taxifolius, 646, 647; tortilis, 151; veracruzensis, 151; viridulus, 151, 196; weirii, 151
Flora additions: to the liverwort flora of Alaska, 63; to the lichen flora of Arizona, 472; California, 469; Colorado, 216; Manitoba, 243; New Mexico, 72; North America, 41, 250; to the moss flora of Arizona, 252; Greenland, 74, 236; Louisiana, 467; North Dakota, 84; South America, 242
Floristics: la brioflora del Macarao, Vene-

zuela, 241; the Hepaticae and Anthocerotae of Florida Parishes, Louisiana, 593, 594*; the Hookeriaceae of Central America, 328; die Lebermoose des Iran und Afghanistans, 48; lichens of the North Cascades Range, Washington, 582, 583*; los Liquenes de Venezuela, 248, 257; the mosses of Costa Rica, 150; the mosses of southeastern Polynesia, 427
Fog effects on lichens, 23
Fontinalis antipyretica, 17-, 447-; novae-angliae, 646
Fontinalis novae-angliae, effects of sudden temperature changes on, 17
Forsstroemia, generic characters, 533*; morphology, 458*
Forsstroemia nitida, 457; trichomitria, 454, 646
Fossils: Epiphyllous moss from the Australian Miocene, 249
Fossombronia, 51, 281; **alaskana** sp. nov., 66, 68*, diagnostic characters, 64*, 65*, 67*; brasiliensis, 596; foveolata, 63-, 596; grossepapillata, 63-; longiseta, 281-; lophoclada, 63-; wondraczekii-longiseta complex, 63-
FREY, W.: Die Lebermoose des Iran und Afghanistans, 48
Frullania eboracensis, marginal gemmae in, 460, 461*
Frullania falciloba, SEM of spores and elaters, 303, 306*, 307*
Frullania asagrayana, 281-, 460-, aubertii, 644; bolanderi, 460-; brittoniae, 460-, 596; campanulata, 460-; dilatata, 52; eboracensis, 596; fragilifolia, 460-; inflata, 413, 556, 596; kunzei, 596; mayebarae, 281-; obcordata, 596; plana, 460-; riparia, 460-; squarrosa, 597; takayuensis, 281-; tamarisci, 52
FULFORD, M.: Recent literature on hepatics, 67, 85; 68, 258; 69, 478
Fumarprotocetraric acid, 128-
Funaria bonplandii, 156; calvescens, 156; epipedostegia, 156; hygrometrica, 6, 172-, 415, 632-, 646; hygrometrica var. calvescens, 156; longiseta, 156; polaris, 175, 195
Funaria hygrometrica, effect of Cu and Zn on growth, 447, 448*; enzymes, 577

GALUN, M. *See* MARTON, K. and M. GALUN
Gametangia, in Splachnaceae, 623
Gametophore development from irradiated spores in Polytrichum commune, 142
Gamma radiation, effects on growth and development of Polytrichum commune, 142, 144*, 145*
Garovaglia obtusata, 437; plicata, 437; tahitense var. brevicuspidata, 437; tahitensis, 437
Gemmae: filamentous in Hepaticae, 57; marginal in Frullania eboracensis, 460, 461*
Genetics and spore markings in liverworts, 324
GEORGE, C. J. *See* SMILEY, D. and C. J. GEORGE
Geothallus tuberosus, 281-
Gibberellins and thallus elongation in Marchantia polymorpha, 33, 34*, 35*, 36*, 37*
GLIME, J. M. and R. E. CARR: Temperature survival of Fontinalis novae-angliae, 17
Glossadelphus laevifolius, 168; longisetus, 168; tahitensis, 441; torrentium, 441; truncatulus, 168
Glyptothecium, 531-
Graphis scripta, 568, 587
GREENBERG, J. *See* MEYER, M. W., J. GREENBERG, and S. TEDESCHI
Greenland, Dicranella riparia new to, 236; Leskeella and Pseudoleskeella in, 601; mosses new to, 74
Grimmia subg. Tricostatae, 233
Grimmia (Coscinodon) **arctolimnia** sp. nov., 230, 231*; calyptrata, 646; contermina, 156; cribrosa, 232; doniana, 447-; laevigata, 647; ovalis, 155; torquata, 74-; trinervis, 233
GROSSMAN, H. H. and C. J. HILLSON: The effect of chronic low-level gamma radiation on Polytrichum commune. I. Growth and development of gametophores from irradiated sporocytes and spores, 142
Groutia, *543*; abietina, *545*
Groutiella apiculata, 159; mucronifolia, 159; schlumbergeri, 159; undosa, 159; wagneriana, 159
Growth effects of ilmenite basalt and volcanic ash on Marchantia polymorpha, 632
Growth of Funaria hygrometrica, 447; of Marchantia polymorpha, 447
GUERKE, W. R.: A floristic study of the Hepaticae and Anthocerotae of the Florida Parishes, Louisiana, 593
Gyalecta erythrozona, 43
Gyalectidium filicinum, 248
Gymnomitrium, 63-
Gymnostomum valeriana, 154
Gyrophoric acid, 128-, 254

Haematomma lapponicum, 587
HALE, M. E. and P. G. PATWARDHAN: Two new species of Hypotrachyna (Lichenes) from India, 637
HANCOCK, J. A. and G. R. BRASSARD: Phenology, sporophyte production, and life history of Buxbaumia aphylla in Newfoundland, Canada, 501
Haplocladium microphyllum, 7, 415, 555
Haplohymenium triste, 165
Haplomitrium gibbsiae, SEM of spores and

elaters, 285*, 286, 287*; TEM and SEM of water conducting elements, 612, 613*, 615*
Haplomitrium hookeri, 236-; rotundifolium, 615*
Haptera, 249*
Harpophyllum aureum, 164
Harrisonia apiculata, 159
HATTORI, S. (review), R. M. Schuster, The Hepaticae and Anthocerotae of North America, East of the Hundredth Meridian, 657
HÉBANT, C. *See* BURR, R. J., B. G. BUTTERFIELD, and C. HÉBANT
Hedwigia ciliata, 6, 74-
Hedwigidium integrifolium, 159
HEGEWALD, E.: Beitrage zur Moosflora Perus. I. Bryomanginia saint-pierrei—neu für Südamerika, 641
Helicophyllum torquatum, 159
Helodium blandowii, 12
Hemiragis aurea, 164, 356
Heppia, 469-; **echinulata** sp. nov., 239*, 240*; lutosa, 214, 587
Heppiaceae, a new type of soredium, 208
Herberta alpina, SEM of spores and elaters, 288, 289*
Herberta divergens, 644
HERMANN, F. J. and H. ROBINSON: Additions to the bryophyte flora of Bolivia, 643
Heterocladium papillosum, *608*
Heterodermia hypoleuca, 244, 414; obscurata, 257; tremulans, 257; 569
HICKS, M. L.: Marginal gemmae in Frullania eboracensis, 460
HILLSON, C. J. *See* GROSSMAN, H. H. and C. J. HILLSON
Himantocladium cyclophyllum, 438; implanum, 438
HOFFMAN, G. R.: Growth stimulation of Marchantia polymorpha from ilmenite basalt and volcanic ash, 632
HOLLENSEN, R. H. (review), J. H. Troughton and F. B. Sampson, Plants. A scanning electron microscope survey, 497
HOLLINGSWORTH, P. J. *See* TAYLOR, J., P. J. HOLLINGSWORTH, and W. C. BIGELOW
Holoblepharum, 438
Holomitrium arboreum, 153; cylindraceum, 431; flexuosum, 153; longifolium, 153; standleyi, 153; terebellatum, 153; vaginatum, 431; williamsii, 153
Homalia glabella, 162; pseudo-exigua, 438
Homaliodendron decompositum, 162; exiguum, 438; flabellatum, 438
Homalotheciella subcapillata, 646
Homalothecium leskeoides, 165
Homalothecium sp., soluble sugars in, 471*
Homosekikaic acid, 617
Hookeriaceae of Central America, 328
Hookeria sect. Callicostella subsect. Thamniopsis, *401*
Hookeria acutifolia, 163, 357; bromeliae, *397*; crispa, 163, *360*; diffusa, *361*; falcata, 163; falcatula, 163; flexifolia, *379*; haplociliata, *383*; nicaraguensis, *347*; nitida, 398; oblongifolia, 438; oerstediana, 163, *339*; pendula, *402*; subfalcata, *369*; taylorii, 370; tenuifolia, *397*; variabilis, *370*; wrightii, *372*
Hookeriopsis sect. Thamniopsis, *401*
Hookeriopsis angustiretis, 358, 359*; callicostelloides, 359*, 360; crispa, 163, 359*, 360; cruegeriana, 361; diffusa, 361, 362*; dimorpha, 363; falcata, 163, 363; falcatula, 163, 363; guadalupensis, 363; gautemalensis, 362*, 363; heteroica, 468; incurva, 364; laevinervis, 163, 362*, 364; obtusifolia, 163, 365, 366*; panamensis, 365, 366*; pendula, *402*; standleyi, 367, 368*; subfalcata, 163, 368*, 369; taylorii, 370; undata, 370; variabilis, 163, 368*, 370
Hormocysts, formation, 210
Hydrocryphaea, 531-
Hydroids, 616
Hylocomium splendens, 12
Hymenophyton flabellatum, TEM and SEM of water conducting elements, 612, 614*, 615*
Hymenostomum breutelii, 154; mexicanum, 154
Hyophila beruensis, 433; subcontermina, 155; tortula, 155
Hypnella jamesii, 168; pilifera, 164, 371; wrightii, 372*
Hypnodendron comosum, 435; junghuhnii, 434; reinwardtii, 434; samoanum, 434; spininervium, 434; tahiticum, 435; vescoanum, 434
Hypnum amabile, 168; apertum var. condensatum, 442; bryifolium, 442; calpaecarpum, 442; catenulatum, *605*; chamissonis, 442; contorte-operculatum, 166; cupressiforme, 415; cupressiforme var. cupressiforme, 6; cupressiforme var. lacunosum, 168; fuscescens, 441; inflectens, 442; lindbergii, 8; loxocarpum, 442; malacodes, 168; mirabile, 168; molluscoides, 442; paradoxum, 440; pickeringii, 441; plumulosum, 440; polypterum, 168; polypterum var. robustum, 168; pratense, 7; pseudoprotense, 165; pycnophyllum, 441; revolutum, 6, 601-; stereodon tectorum, *607*; tectorum, *606*, *607*
Hypogymnia atrofusca, 473; austerodes, 218; bitteriana, 218; enteromorpha, 587; intes-

tiniformis, 587; physodes, 569, 587; tubulosa, 218; vittata, 587
Hypoprotocetraric acid, 72-
Hypopterygium arbusculosum, 439; debile, 439; filiculaeforme, 439; nadeaudianum, 439; rigidulum, 164; spectabile, 436; tahitense, 439; tamariscinum, 164; tomentosum, 159
Hypotrachyna **awasthii** sp. nov., 637, 638*; **dodapetta** sp. nov., 637, 638*; kingii, 639; rhabdiformis, 637; thryptica, 637

Icmadophila ericetorum, 587
INOUE, H. *See* STEERE, W. C. and H. INOUE
International Association of Bryologists, 105*
Iran, die Lebermoose, 48
IRELAND, R. R.: Neomacounia, a new genus in the Neckeraceae (Musci) from Canada, 453
Isodrepanium lentulum, 164, 373
Isopterygiopsis, 75
Isopterygium albescens, 442; albescens var. applanata, 442; argyrocladum, 442; chrismarii, 644; cylindricarpum, 168; diminutivum, 166; elegans, 74-; fecundum, 166; integrifolium, 166; minutirameum, 442; minutirameum var. brevifolia, 442; miradoricum, 167; muellerianum, 74*, 75; perpallidum, 442; poasense, 167; pusillum, 167; semicostatum, 167; subtenerrimum, 167; subtrichopelma, 167; tenerum, 167, 646; trichopelma, 167
Isothecium bonplandii, 165
Israel, a new Heppia from the Arava Valley, 239

Jaegerina scariosa, 160
Jamesoniella autumnalis, 8, 597
JOHNSEN, A. B.: Tayloria splachnoides in Arizona, 636
JONESCU, M. E. *See* SHEARD, J. W. & M. E. JONESCU
Jungermannia gracillima, 597
Jungermannia hodgsoniae, SEM of spores and elaters, 295, 296*

KALAITZAKIS, J. A. *See* MARGARIS, N. S. and J. KALAITZAKIS; and N. S. MARGARIS: Quantitative analysis of soluble sugars from two lichen species, 465
KITAGAWA, N. and T. KODAMA: A remarkable new species of Acromastigum (Hepaticae) with septate rhizoids and filamentous gammae, 57
KODAMA, T. *See* KITAGAWA, N. and T. KODAMA
Koerberia sonomensis, 219
KROG, H.: Parmelia ultralucens, a new lichen species in subgenus Amphigymnia, 253

Lasallia papulosa, 179; pennsylvanica, 181
Latin America: Bolivia, bryophyte additions, 643; Bryomanginia saint-pierrei, new to, 641; Central America, Hookeriaceae of, 328; Costa Rica, mosses from, 150; Placomaronea candelarioides in, 463; Syrrhopodon quintasii, new to, 242; Venezuela, Brioflora de, 241; Venezuela, Liquenes de, 248, 257
LAZARENKO, A. S.: Some considerations on the nature and behavior of the relict moss, Desmatodon randii, 474
Lecania dimera, 244, 414, 587; syringea, 516
Lecanora, 82-; sect. Aspicilia, 128-
Lecanora alpina, 587; atra, 219, 257; cadubriae, 219; caesiocinerea, 587; calcarea, 587; cenisia, 587; chloropolia, 219; chrysoleuca, 587; cinereofusca, 587; cinereorufescens, 216; coilocarpa, 516, 587; epibryon, 42, 587; frustulosa, 587; gibbosula, 587; granatina, 219; hageni, 414, 587; hypoptoides, 516; impudens, 244, 414; laevata, 587; melanophthalma, 587; mougeotioides, *132*; muralis, 587; pacifica, 587; perradiata, 140; piniperda, 473, 567; polytropa, 587; pruinosa, 42; rupicola, 587; sorediza, 140; **stellata** comb. nov., 137*, 140; straminea, 42; straminea var. oreina, *132*; subfusca, 587; supertegens, 587; umbrosa, 43; varia, 587
Lecanoric acid, 72-
Lecidea, 82-; aeruginosa, 219; albocaerulescens, 587; alpestris, 44; anthracophila, 473; armeniaca, 473; atrobrunnea, 587; auriculata, 587; berengeriana, 473, 587; caesioatra, 587; cf. cinerata, 587; circumnigrata var. circumnigrata, 220; circumnigrata var. reagens, 219; confluens, 587; crustulata, 588; cyathoides, 588; demissa, 588; demonum, *219*; elata, 473; elegantior, 473; furvonigrans, 588; fuscescens, 588; glaucophaea, 588; globifera, 588; glomerulosa, 414, 588; granulosa, 217, 588; lactea, 216, 219; lapicida, 588; lurida, 588; lyngei, 473; macrocarpa, 588; marginata, 588; paddensis, 588; plana, 588; populina, 83; rhaetica, 220; rubiformis, 588; rufonigra, 588; sanguineoatra, 588; silacea, 220; speirea, 220, 588; symmicta, 567; tesselina, 473; tessellata, 588; tornoensis, *220*, 588; tuberculata, *46*; uliginosa, 588; vernalis, 568
Lecidella bullata, 44; carpathica, 45; elaeochroma, 588; goniophila, 473; incongruella, 45; stigmatea, 588
Leiomela bartramioides, 158
Leiomitrium standleyi, 158
Lejeunea cavifolia, 52; cladogyna, 597; flava, 597; laetevirens, 597; ulcina, 597

Lempholemma, 210
Lepicolea attenuata, SEM of spores and elaters, 288, 290*, 292*
Lepidolaena clavigera, SEM of spores and elaters, 291, 293*, 294*
Lepidopilidium divaricatum, 375; portoricense, 163, 374; subdivaricatum, 163, 372*, 374
Lepidopilum apiculatum, 164, 377, 378*; apophysatum, 379; biductulosum, 378*, 379; brevipes, 164, 380; carneum, 164, 380; contiguum, 164, 378*, 381; cubense, 164, 381; diaphanum, 381; falcatulum, 382; flexifolium, 379; floresianum, 164, 382, 383*; haplociliatum, 164, 383*; laetenitens, 164; livens, 164, 384, 386*; mohrianum, 385; muelleri, 385; platyphyllum, 164, 385, 386*; polytrichoides, 164, 387; polytrichoides var. costaricense, 164; pumilum, 388, 389*; radicale, 164, 388; rotundifolium, 164, 389*, 390; scabrisetum, 164, 390; semilaeve, 164, 391, 392*; skutchii, 164, 391, 392*; stolonaceum, 393; subdivaricatum, 163, *374*; subenerve, 164; subtortifolium, 393, 394*; tortifolium, 164, 394; wallisii, 346
Lepidozia spinosissima, SEM of spores and elaters, 295, 297*
Lepp, N. W. *See* Coombes, A. J. and N. W. Lepp
Lepraria, 567; membranacea, 588; neglecta, 588
Leptobryum pyriforme, 7, 74-, 172-
Leptochidium albociliatum, 220
Leptodictyum riparium, 6, 415; trichopodium, 12, 415, 556
Leptodon abietinus, *545*; circinalis, *545*; circinatus, *545*; circinnatus, *546*; nitidus, *457*; smithii, 454, 537; trichomitrion, 454
Leptodontis, *545*
Leptodontium araucarietii, 644; brachyphyllum, 644; capituligerum, 155; exasperatum var. flagellaceum, 155; excelsum, 155; filescens, 155; filescens var. denticulatum, 155; filicola, 155; flexifolium, 155; longicaule, 155; motelayi, 155; orcuttii, 155; subgracile, 155; sulphureum, 155; sulphureum var. flagellaceum, 155; sulphureum var. panamense, 155; syntrichioides, 155; ulocalyx, 155; valeriana, 154; viticulosoides, 644; viticulosoides var. exasperatum, 155; wallisii, 155
Leptogium austroamericanum, 473; lichenoides, 588; marginellum, 257; tenuissimum, 588
Leptostomum sp., 434; macrocarpum, 434
Leptotheca costaricensis, 157
Leptotrichum costaricense, 153
Lepyrodon tomentosus, 160; tomentosus var. latifolius, 160
Leskea, 442; bulbifera, *602*; catenulata, *605*; cyrtophylla, *609*; extensa, *602*; montanae, *603*; nervosa, *602*; norvegica, *602*; obscura, 415; obscura/polycarpa, 556; papillosa, *608*; polycarpa, 415; tectorum, *606*; wollei, *606*
Leskeella nervosa, 602*, 603*; tectorum, *606*
Leskeodon andicola, 395; cubensis, 395; longipilus, 395; mexicanus, 396; pusillus, 163, 396
Letharia vulpina, 588; diurnal cycle of water content, 25*; effects of humidity on thallus water content, 27*; water relations, 23, 27*, 28*
Leucobryum albicans, 153; albidum, 154, 646; antillarum, 154; costa-ricense, 154; crispum, 154; crispus, 241; giganteum, 154; longifolium, 154; martianum, 154; mittenii, 154; polakowskyi, 154; scalare, 431
Leucodon, generic characters, 523*
Leucodon curvirostris, 160; domingensis, 160; julaceus, 646; pacificus, 436
Leucodontaceae, generic data matrix, 533*; revision, 531; subfam. Alsioideae, revision, 531, 538; subfam. Antitrichioideae, *536*; subfam. **Felipponoideae** subfam. nov., 535; subfam. **Forsstroemioideae** subfam. nov., 535; subfam. Leucodontoideae, 536; subfam. Pterogonioideae, *536*; tribe Alsieae, *538*; tribe Antitrichieae, *536*; tribe **Dozyeae** tribe nov., 536; tribe Leucodontae, 536; tribe Pterogonieae, 536
Leucodontopsis, generic characters, 533*
Leucodontopsis floridana, 160; geniculata, 160, 242-; plicata, 160
Leucolejeunea clypeata, 597; conchifolia, 597; unciloba, 597
Leucoloma crugerianum, 153; serrulatum, 153; subimmarginatum, 153; molle, 431; tortellum, 153
Leucomium aneurodictyon, 439; costaricense, 164; debile, 439; lignicola, 164
Leucophanes albescens, 431; blumii, 432; calymperatum, 154; fragile, 432; nukahivense, 431; octoblepharioides, 431; prasiophyllum, 431; tahiticum, 432
Lewinsky, J.: Some new or noteworthy mosses from west Greenland, 74; The genera Leskeella and Pseudoleskeella in Greenland, 601
Lichen chemistry, 203
Lichenic acid, 220
Lichens: air pollution, 561; distribution in west central Canada, multivariate analysis, 514, 519*, 523*; in the Shawangunk

Mountains of New York, 179; physical condition and distribution on four tree species, 568*
Lichexanthone, 251, 253
Lichina pygmaea, sugars in, 466
Lindigia aciculata, 161
Literature on hepatics, 67, 85; 68, 258; 69, 478
Literature on lichens, 87, 95; 88, 267; 89, 488; 90, 648
Literature on mosses, 97, 88; 98, 262, 99, 481
Lobaria linita, 588; oregana, 588; pulmonaria, 588; quercizans, 569; scrobiculata, 212, 588
LONGTON, R. E.: Microclimate and biomass in communities of the Bryum association on Ross Island, continental Antarctica, 109
Lopadium puiggarii, 248
Lophocolea bidentata, 8, 597; heterophylla, 7, 51, 57-, 281-, 597; horikawana, 57-; martiana, 597; minor, 8, 51, 636-
Lopholejeunea muelleriana, 597
Lophozia badensis, 51; bicrenata, 632-; longidens, 555; turbinata, 51; ventricosa, 555; wenzelii, 74-
Lopidium trichocladulum, 439
Louisiana, Hepaticae and Anthocerotae of the Florida Parishes, 593, 594*; mosses, 467
Loxodin, 250
Lunularia cruciata, 50, 281-
Lysosomal degeneration in water conducting strands, 615

Macrocoma hymenostomum, 159; sullivantii, 159
Macromitrium apiculatum, 159; barbense, 158; barbense var. jamaicense, 158; **bescherellei** nom. nov., 435; bistratosum, 435; cacuminicola, 436; cirrosum, 158; crenulatum, 158; didymodon, 159; durandii, 159; eurymitrium, 435; fragile, 159; fulgescens, 158; fuscescens, 158; fusco-aureum, 158; greenmanii, 158; guatemalense, 158; hirtellum, 159; homalacron, 158; incurvifolium, 436; lamprocarpum, 159; longifolium, 158; longifolium var. viridissimum, 158; mammillosum, 158; menziesii, 436; mexicana, 159; mucronifolium, 159; palmense, 158; paridis, 436; parvirete, 159; pentastichum, 159; podocarpi, 159; punctatum, 159; rapaense, 436; reflexifolium, 159; reinwardtii, 436; richardii, 159; ruginosum, 436; savatieri, 436; schlumbergeri, 159; scleropelma, 158; serrulatum, 159; standleyi, 159; standleyi var. subundulatum, 159; st.-johnii, 436; stratosum, 159; subcirrhosum, 159; suberosulum, 436; subtile, 436; tahitense, 435; tonduzii, 159; verrucosum, 159; werckleanum, 158; williamsii, 159
Madotheca levigata, 52
MAGILL, R. E.: Notes on Texas mosses III, 645; and F. SEABURY and D. M. J. MUELLER: Evaluation of the critical point drying technique and its application in systematic studies of mosses, 628
Malate dehydrogenase in Funaria hygrometrica, 577, 578*, 579*
Manitoba, Bird's Hill Provincial Park, map, 2*; bryophytes, association table, 7*; corticolous bryophytes, 551; ecological studies of bryophytes of coniferous forests, 1; lichens, additions, 243; lichens and bryophytes, effects of air pollution, 405
Mannia androgyna, 49; fragrans, 49
MANUEL, M. G.: A revised classification of the Leucodontaceae and a revision of the subfamily Alsioideae, 531
MARAVOLO, N. C. *See* MELSTROM, C. E., N. C. MARAVOLO, and J. R. STROEMER
Marchantia berteroana, SEM of spores and elaters, 317, 319*
Marchantia domingensis, 597; paleacea, 50; polymorpha, 7, 50, 281-
Marchantia polymorpha, effect of Cu and Zn on growth, 447, 448*, 450*; effects of gibberellins on growth, 33, 38*, 39*; effects of ilmenite basalt and volcanic ash on growth, 632
MARGARIS, N. S.: Free amino acid pools in Cladonia pyxidata and Peltigera sp., 77; and J. KALAITZAKIS: Soluble sugars from some Greek mosses, 470; *see* KALAITZAKIS, J. A. and N. S. MARGARIS
Maronella, 220
MARTON, K. and M. GALUN: A new species of Heppia from the Arava Valley, Israel, 239
Massalongia carnosa, 588
Mastigolejeunea auriculata, 597
Mazosia rotula, 248
Megalospora versicolor, 257
Meiotheciopsis lageniformis, 167
Meiothecium hamatulum, 441; intextum, 441; rechingeri, 441; serrulatum, 441; tenerum, 167
Melanaspicilia, 128-
MELSTROM, C. E., N. C. MARAVOLO, and J. R. STROEMER: Endogenous gibberellins in Marchantia polymorpha and their possible physiological role in thallus elongation and orthogeotropic growth, 33
Menegazzia terebrata, 588
Merceya agoyonensis, 154; ligulata, 154, 233
Mesonodon onustus, 166
Mesoptychia sahlbergii, 195-

Meteoriopsis consimilis, 161; patula, 161; recurvifolia, 161; remotifolia, 161
Meteorium, 437; patula, 161; undulifolium, 161
Method to measure water uptake in lichens, 25
Metzgeria conjugata, 51; filicina, 647; furcata, 50; myriopoda, 597
Metzlerella costaricensis, 153; longiseta, 153
Mexico, Archidium, new genus to, 639
MEYER, M. W., J. GREENBERG, and S. TEDESCHI: Enzymes of the moss Funaria hygrometrica II: The isoenzymes of malate dehydrogenase, 577
Micarea, 220; confusula, 82-; excimia, 82-; gelatinosa, 82-; lithinella, 82-; **populina** comb. nov., 83; **tuberculata** comb. nov., 46
Microclimate: of Bryum associations, 109; at Cape Armitage, Antarctica, 112*, 113*, 114*, 115*, 116*, 117*, 118*, 119*, 120*, 121*; determination, 111
Microdus barbensis, 152; kunzeanus, 152; longirostris, 152; picquenotii, 431
Microenvironment of Ramalina menziesii, 23
Microlepidozia sylvatica, 597
Micronutrients: in ilmenite basalt, 633*, 634*; in volcanic ash, 634*
Microthamnium, 442; atro-viride, 168; elegantulum, 169; incompletum, 169; isopterygioides, 169; langsdorffii, 169; laxulum, 169; lehmanni, 169; minisculifolium, 169; reptans, 169; rostratulum, 169; sordidum, 169; thelistegum, 168; turckheimii, 169
Mielichhoferia graciliseta, 156; macrodonta, 156; minutifolia, 156; nitida, 233; practicola, 156
MILLER, N. G. (review) H. Crum, Mosses of the Great Lakes Forest, 100
Miocene, epiphyllous moss, 249
Mittenothamnium atro-viride, 168; diminutivum, 168, 241; elegantulum, 169; incompletum, 169; isopterygioides, 169; langsdorffii, 169; laxulum, 169; lehmannii, 169; macroblepharum, 442; minisculifolium, 169; reptans, 169, 241; rostratum, 169; sordidum, 169
Mniadelphus montagneanus, 439
Mniodendron, 435
Mnium affine, 646; cuspidatum, 8, 415, 556; cuspidatum union, 8; drummondii, 7; rostratum, 157, 434; rugicum, 8, 415; rugicum-Thuidium recognitum union, 8; spiniforme, 157; spinosum, 195-; spinulosum, 6; spinulosum in North America, 79, 80*; undulatum, 623-
Moenkemeyera rarotongae, 430
Moerckia flotoviana, 615*
MOLENAAR, J. G. DE: Dicranella riparia new to Greenland, 236
Monoclea forsteri, SEM of spores and elaters, 311*, 312
Morinia ehrenbergiana, 155
mucilage hairs, 623-
MUELLER, D. M. J. *See* MAGILL, R. E., F. SEABURY, and D. M. J. MUELLER
Multivariate analysis of lichen distributions and species associations, 514, 520
MURRAY, B. M. *See* STEERE, W. C. and B. M. MURRAY
Mycoblastus sanguinarius, 588; **tornoensis** comb. nov., 220
Myurella julacea, 6
Myurium, 531-

Nadeaudia schistostegiella, 434
Naiadita lanceolata, archegonia, 627
Nardia, 63-
NASH, T. H.: New and additional Xanthoparmelia from New Mexico, 72; Two new species of Xanthoparmelia with diffractaic acid, 234; (review) V. Ahmadjian and M. E. Hale (editors), The Lichens, 661; and W. A. WEBER: Additions to the lichen flora of Arizona II, 472
Neckera, 458; abietina, *545*; angustifrons, *373*; australasica, 438; biductulosa, *379*; californica, *539*; chlorocaulis, 162; cladorrhizans, 440; cylindricaule, 166; disticha, 162; falcifolia, 164; graeffeana, 438; graeffeana var. tahitensis, 438; implana, 438; lepineana, 438; macounii, *457*; madagascariensis, 441; nigrescens, 161; oerstediana, 161; pennata, 413, 428; rigida, 161; undulata, 438; wercklei, 162
Neckeropsis disticha, 162; foveolata, 162; lepineana, 438; undulata, 162
Neohodgsonia mirabilis, SEM of spores and elaters, 320, 321*, 322*
Neohypnella diversifolia, 164, 396; mucronifolia, 164
Neomacounia gen. nov., 454, morphology, 458*; **nitida** comb. nov., 455*, 456*, 457
Nephroma bellum, 588; helveticum, 571, 588; laevigatum, 588; parile, 212, 213*, 588; resupinatum, 473, 588
Neuropogon lambii, 588
NEWBERRY, G.: The influence of a sulfate-process paper mill on corticolous lichens, 561
New Mexico, new and additional Xanthoparmelia, 72
News and Notes, 105, 280, 499, 663
New York, lichen decline in the Shawangunk Mountains, 179, 180*, 181*, 182*, 183*, 184*, 185*, 186*
Norlobaridone, 250, 255

Normandina pulchella, 257, 469
Norstictic acid, 128-, 220, 637
North America: additions to the lichen flora of, 41; Dimelaena of, 128; distribution of Bryum wrightii in arctic and boreal regions of, 172; Mnium spinosum in, 79; a new lichen in, 250; the Ramalina intermedia complex of, 617
North Dakota mosses, 84
Northwest Territories, a new species of Grimmia from, 230
Notothylas breutelii, 596
Nowellia curvifolia, 281-

Obtusatic acid, 619
Ochrobryum gardneri, 154; obtusifolium, 154
Ochrolechia androgyna, 473, 589; frigida, 42; oregonensis, 589; pallescens, 569, 589; cf. parella, 589; tartarea, 589; upsaliensis, 589
Octoblepharum albidum, 154, 432, 593-; cocuiense, 154; erectifolium, 154; longifolium, 432; pulvinatum, 154
Odontoschisma denudatum, 596; prostratum, 596
Oedicladium, 532
Oligotrichum aligerum, 169; erosum, 169
Omphalodiscus decussatus, 589; kraschenninikovii, 469; virginis, 589
Oncophorus riparius, 236; wahlenbergii, 6, 74-
Opegrapha filicina, 248; pulicaris, 244, 414, 589
Oreas martiana, 79-
Orthodicranum flagellare, 153
Orthodontium gracile, 156; pellucens, 156
Orthogeotropic growth in Marchantia polymorpha, 33
Orthorrhynchium cylindricum, 437
Orthostichidium pentagonum, 160
Orthostichopsis tetragona, 160
Orthotrichum, 601-; obtusifolium, 6, 413; pumilum, 6, 413, 556; speciosum, 13, 413
Osorio, H. S.: Placomaronea candelarioides: Ampliación de su área de distribución geográfica, 463
Oxylate crystals, 128-
Oxyrrhynchium rugosipes, 440
Oxystegus cylindricus, 155

Pallavicinia lyellii, 597
Pannaria granatina, 219; pityrea, 212, 213*; rubiginosa, 589
Papillaria aongstroemii, 437; appressa, 161; appressa var. illecebra, 161; deppei, 161; helictophylla, 437; imponderosa, 161; nigrescens, 161; oerstediana, 161; tahitensis, 437; viridula, 160
Paraleucobryum albicans, 153
Parmelia subg. Amphigymnia, 234-, 251, 253; subg. Imbricaria, 253-; subg. Hypotrachyna, 253-; subg. Parmelia, 253-; subg. Parmelia sect. Irregulares, 234-, 250-; subg. Xanthoparmelia, 234
Parmelia, 82-; **ajoensis** sp. nov., 234, 235*; albertana, 414; alpicola, 589; aspera, 473; aurulenta, 567; bolliana, 244, 414, 567; caperata, 414, 567; commensurata, 250; conspersa, 221, 589; crinita, 256; dierythra, 73, 235; diffractaica, 250; dilatata, 257; dissecta, 220; dissensa, 73; farinacea, 218; flaventior, 414, 567; formosana, 253-; galbina, 567; infumata, 589; **joranadia** sp. nov., 72; koyaensis, 256; kurokawae, 220, 235; lecanorica, 72-; lineola, 235; monumentalis, 253; multispora, 589; panniformis, 589; perlata, 589; psoromifera, 235; reticulata, 221, 250, 253-; rudecta, 567; saxatilis, 567, 589; simulans, 250; sorediosa, 221; subargentifera, 244, 414; subaurifera, 567; subcrinita, 251, 253; subdecipiens, 221; subisidiosa, 256; subrudecta, 569; subtinctoria, 255; sulcata, 414, 567, 589; tasmanica, 73; tinctina, 73; **tucsonensis** sp. nov., 234, 235*; ulophyllodes, 245, 414, 567; **ultralucens** sp. nov., 253, 254*, 255*; wallichiana, 256, 637; weberi, 235; worcesteri, 72-; wyomingica, 473; xanthina, 257
Parmeliella cyanolepra, 589; praetermissa, 589
Parmeliopsis ambigua, 589; hyperopta, 589
Patwardhan, P. G. *See* Hale, M. E. and P. G. Patwardhan
Pellia endiviifolia, 51; epiphylla, 33-, 51, 597; fabroniana, 597
Peltigera sp., soluble sugars in, 465*, 466*, 467*
Peltigera aphthosa, 589; canina, 589; canina var. spuria, 212, 213*; collina, 589; elisabethae, 221; lepidophora, 589; malacea, 589; membranacea, 589; polydactyla, 589; rufescens, 589; spuria, 589; venosa, 473, 589
Peltula bolanderi, 210, 211*, 469-; clavata, 212; euploca, 128-, 209*, 210, 211*, 213*, 469-; placodizans, 210, 211*; zahlbruckneri, 470
Pertusaria amara, 589; glaucomela, 589; leioplaca, 589; multipuncta, 589; octomela, 221; pertusa, 569, 589; saximontana, 221; sommerfeltii, 221, 473; velata, 567
Peru, Bryomanginia saint-pierrei, neu für, 641; Placomaronea candelarioides in, 463
Phaeoceros laevis, 596; laevis subsp. laevis, 49
Phenology of Buxbaumia aphylla, 501; of Desmatodon randii, 474
Philonotis, 109-; crassinervia, 158; glaucescens, 158; hastata, 435; jardini, 435; longiseta, 158; marchica, 468; nanodendra, 158;

runcinata, 435; sphaericarpa, 158; st.-johnii, 435; tahitensis, 435; tenella, 158; tomentella, 238; uncinata, 158, 468; vescoana, 435
Philonotula jardini, 435
Philophyllum bromeliae, *397*; tenuifolium, 394*, 397
Phyllobathellium epiphyllum, 248
Phyllogonium, 438; fulgens, 161; fulgens var. gracile, 161; fulgens var. viride, 161; viscosum, 162
Physcia, 128-, 208-; adscendens, 414, 516, 568; aipolia, 414, 516, 567, 589; caesia, 589; cascadensis, 590; ciliata, 414, 516; dubia, 590; elaeina, 567; grisea, 567; hassei, 128-; hirsuta, 82-; leptalea, 516; millegrana, 245, 414, 567; orbicularis, 212, 213*, 414, 516, 567; orbicularis fo. albociliata, 567; orbicularis fo. rubropulchra, 567; stellaris, 414, 516, 567, 590; tenella, 414; teretiuscula, 590; tribacia, 590; wainioi, 473
Physcomitrium pyriforme, 6, 646
Physconia enteroxantha, 221; grisea, 414; muscigena, 42, 590; pulverulenta, 590
Pilophoron acicularis, 590; cereolus, 590; clavatus, 590
Pilopogon gracilis, 153; gracilis var. pittieri, 153
Pilosium longisetulum, 166
Pilotrichella flexilis, 161; hexasticha, 161; isoclada, 161; longipila, 161; pentasticha, 161; pulchella, 161; quinquefaria, 160; rigida, 161; tenuinervis, 161; tetragona, 160; tonduzii, 161
Pilotrichopsis, generic characters, 533*
Pilotrichum abietinum, *545*; amazonum, 163; andersonii, 162; asperifolium, 162; biductulosum, 379; bipinnatum, 163; compositum, 163; evanescens, 163; fendleri, 163; mucronatum, 163; mucronatum var. elongatum, 163; pallidum, 163; ramosissimum, 163; tonduzii, 163; undulatum, 162
Pinnatella elegantissima, 438; kuehliana, 438; minuta, 162
Pirea mariae, 161
Pireella angustifolium, 160; cymbifolia, 160; mariae, 161; pohlii, 161
Placomaronea candelarioides, 463, 464*; candelarioides var. lacinulata, 463
Placopsis gelida, 590
Plagiochasma, 49; rupestre, 644
Plagiochila aerea, 644; asplenioides var. porelloides, 51; dubia, 597; horrida, 644; ludoviciana, 597; miradorensis, 597; paradoxa, *60*; undata, 597; verruculosa, 644
Plagiomnium rostratum, 434; rugicum, 74-
Plagiopus oederi, 77
Plagiothecium denticulatum, 6, 166, 636-; piliferum, 74-; planissimum, 166, 242
Plasmodesmata, 616
Platismatia glauca, 590; herrei, 590; lacunosa, 590; norvegica, 590
Platydictya subtile, 7
Platygyrium repens, 7, 647
Platyhypnidium aquaticum, 165; obscurum, 440
Platyhypnidium riparioides, soluble sugars in, 471*
Pleuropus bonplandii, 165
Pleurothotrichum cryptocarpum, 169; nukahivense, 436
Pleurozium schreberi, 8, 166
Pogonatum aloides, 142-; alpiniforme, 170; andinum, 169; barbanum, 169; confertidens, 169; consobrinum, 169; convolutum, 169; graeffeanum, 443; hamatifolium, 170; laterale, 443; liebmannianum, 170; octangulare, 170; pennsylvanicum, 646; pittieri, 170; purpurascens, 644; robustum, 170; tahitense, 443; tortile, 170
Pohlia, 109-; commutata, 236-; cruda, 156; flexuosa, 156; integridens, 156; nutans, 6, 33-; papillosa, 156; peracuminata, 156; spectabilis, 156
Polyblastia bryophila, 42, 221; theleodes, 222
Polychidium, 220
Polygraph analysis of lichen distributions and SO_2, 564*, 565*
Polynesia, mosses of southeastern, 427
Polyploidy, in Desmatodon randii, 476
Polysaccharides in liverworts, 615
Polytrichadelphus costaricensis, 170; nukahivensis, 443; peruvianus, 170
Polytrichastrum alpiniforme, 170
Polytrichum commune, effects of gamma radiation on, 142, 146*
Polytrichum antillarum, 170; commune, 644; convolutum var. cirrhatum, 443; juniperinum, 6, 170; octangulare, 170; strictum, 109-; tortile, 170
Porella arboris-vitae, 52; cordaeana, 52; decurrens, 53; pinnata, 597; platyphylla, 413, 597; trigonifolia, 53; ulophylla, 281-
Porella elegantula, SEM of spores and elaters, 299, 305*
Porina emitatrix, 248; epiphylla, 248; nitidula, 248; phyllogena, 248; platypoda, 248; rubentior, 248; rufula, 248; tetramera, 248; thaxteri, 248; verruculosa, 248
Porothamnium crassipes, 162; neckeraeforme, 162
Porotrichodendron substolonaceum, 162; superbum, 162
Porotrichum cobanense, 162; crassipes, 162; decompositum, 162; elegantissimum, 438;

insularum, 162; longirostre, 162; neckeraeforme, 162; pittieri, 162; plagiorhynchum, 162; plicatulum, 162; plumosum, 162; stolonaceum, 162; substolonaceum, 162
Pottia contermina, 155; randii, *474*
Preissia quadrata, 50
Prionodon densus, 160; dichotomus, 160; fusco-lutescens, 160; longissimus, 160; luteovirens, 160; luteovirens var. crispatulus, 160; luteovirens var. mexicana, 160
Protocetraric acid, 619
Protolichesterinic acid, 72-
Protonemata, with haptera, 249*
Pseudocryphaea, generic characters, 533*
Pseudocryphaea flagellifera, 160
Pseudocyphellaria anthraspis, 590
Pseudoleptogium albociliatum, *220*
Pseudoleskea catenulata, *605*; malacoclada, *606*; papillosa, *608*; tectorum, *606*; wollei, *606*
Pseudoleskeella catenulata, 601, 604*; catenulata var. acuminata, 606; nervosa, *603*; papillosa, 608, 609*, 610*; tectorum, 606, 607*, 608*, tectorum subsp. **cyrtophylla** comb. nov., 609
Pseudosymblepharis bartramii, 155; schimperiana, 155
Psoroma hypnorum, 473, 590
Psoromic acid, 41-, 221
Pterigynandrum catenulatum, *604*; filiforme, 74-; filiforme var. mexicanum, 166; nervosum, *602*
Pterobryon angustifolium, 160; densum, 161
Pterobryopsis mexicana, 160
Pterobryum, 437
Pterogonidium pulchellum, 167
Pterogoniella, 441
Pterogonium, generic characters, 533*
Pterogonium catenulatum, *604*; nervosum, *602*; tectorum, *607*
Pterygoneurum, 63-; arcticum, 63-
Ptilidium pulcherrimum, 8, 281-
Ptilium crista-castrensis, 6
Ptychomnion aciculare, 437
Puiggariella aurifolia, 169
Pylaisiella polyantha, 9, 415, 556
Pyxine, 253-; sorediata, 569

Quantitative sampling of bryophyte stands, 3

Raciborskiella prasina, 248; janeirensis, 248
Racomitrium canescens, 74-; crispulum, 156; lanuginosum, 74-; papeetense, 433
Racopilum convolutaceum, 436; cuspidigerum, 436; microphyllum, 436; pacificum, 436; pacificum var. tahitense, 436; spectabile, 436; spectabile var. spinosum, 436; tomentosum, 159, 241
Radula australis, 597; complanata, 7, 51, 281-; jamesonii, 644; lindbergiana, 52; mollis, 597; obconica, 597; sullivantii, 597
Ramalina, 189; capitata, 619; crinata, *191*; farinacea, 590, 617; farinacea var. bolcana, *619*; fastigiata, 567; geniculata, 590; inflata, 590; intermedia, 617, 618*, 620*; maciformis, 23-; melanothrix, *194*; menziesii in California, 24*; minuscula subsp. intermedia, *619*; obtusata, 619; peruviana, 619; **petrina** sp. nov., 621*, 622*; pollinaria, 617; roesleri, 617
Ramalina menziesii: diurnal cycles of water content, 25*; effects of humidity on thallus water content, 27*; morphology of, 23; reticulations, 30*, 31*; water relations, 23, 27*, 28*
Rauia subcatenulata, 165
Rauiella leskeaefolia, 165; subcatenulata, 165
Reboulia hemisphaerica, 49, 281-, 597
Rectolejeunea maxonii, 597
Redfearn, P. L., Jr. (review) D. H. Vitt, A Revision of the Genus Orthotrichum in North America, North of Mexico, 279; (review) W. A. Weber, Guide to the Mosses of Colorado. Keys and Ecological Notes Based on Field and Herbarium Studies, 496
Reese, W. D.: Syrrhopodon quintasii in the Americas, 242; Bryoxiphium norvegicum in Arizona, 252; Notes on Louisiana mosses —VI, 467; (review) W. T. Stearn, Botanical Latin. History, Grammar, Syntax, Terminology, and Vocabulary. 2nd ed., 103; (review) A. Michaux, Flora Boreali-Americani, 495; News and Notes, 105, 280, 499, 663
Relict moss, Desmatodon randii, 474
Reviews: Ahmadjian, V. and M. E. Hale (editors), The Lichens, 661; Birks, H. J. B., Past and Present Vegetation of the Isle of Skye: A Palaeoecological Study, 277; Crum, H., Mosses of the Great Lakes Forest, 100; Degener, O., I. Degener, and H. Hörmann, Mosses of Hawaii, 103; Ferry, B. W., M. S. Baddeley, and D. L. Hawksworth (editors), Air Pollution and Lichens, 101; Flowers, S., Mosses: Utah and the West, 276; Heywood, V. H. (editor), Taxonomy and Ecology. Proceedings of an International Symposium held at the Department of Botany, University of Reading, 660; Inoue, H., Illustrations of Japanese Hepaticae, 659; Martin, W. and J. Child, Lichens of New Zealand, 104; Michaux, A., Flora Boreali-Americani, 495; Nilsson, S. (editor), World Pollen and Spore Flora, 495; Philippi, G., Moosflora

und Moosvegetation des Freeman-Sund-Gebietes (Südost-Spitzbergen), 497; Schuster, R. M., Hepaticae and Anthocerotae of North America, East of the Hundredth Meridian, 657; Stearn, W. T., Botanical Latin. History, Grammar, Syntax, Terminology, and Vocabulary. 2nd ed., 103; Troughton, J. H., and F. B. Sampson, Plants. A Scanning Electron Microscope Survey, 497; Vasari, Y., H. Hyvärinen, and S. Hicks (editors), Climatic Changes in Arctic Areas During the Last Ten-thousand Years. A Symposium Held at Oulanka and Kevo, 4–10 October, 1971, 103; Vitt, D. H., A Revision of the Genus Orthotrichum in North America, North of Mexico, 279; Weber, W. A., Guide to the Mosses of Colorado. Keys and Ecological Notes Based on Field and Herbarium Studies, 496

Reyes, C. R.: Nota adicional acerca del catalogo de los liquenes de Venezuela, 248; and F. Bowers: Contribución a la Brioflora del Macarao, Venezuela I, 241; and A. Skorepa: Contribución á la flora liquenologica del Macarao, Venezuela I, 257

Rhabdoweisia crispata, 74*; fugax, 153; fugax var. tenerrima, 153

Rhacocarpus apiculatus, 159; purpurascens, 160

Rhacopilopsis trinitensis, 169, 242-

Rhamphidium dicranoides, 155; macrostegium, 155

Rhaphidorrhynchium decurvifolium, 241; lindigii, 167; quaylei, 441; subscabrum, 168; subsimplex, 168; xylophilum, 167

Rhaphidostegium barnesii var. costaricensis, 167; caespitosum, 167; caespitosum var. galipense, 167; cuspidiferum, 167; dimorpha, *167*; harpidioides, *167*; loxense, 167; loxense var. cuspidatum, 167; oblique-rostratum var. drepanioides, 167; pickeringii, 441; simulans, *168*; subcespitosum, *168*; tonduzii, *168*

Rhaphidostichum brevisetum, 441

Rhegmatodon rufus, 440

Rhizocarpon disporum, 590; distinctum, 590; geographicum, 590; hochstetteri, 590; macrosporum, 590; obscuratum, 222, 590; cf. jemtlandicum, 590; permodestum, 46; tetramerum, 222

Rhizogonium lindigii, 157; mnioides, 157; setosum, 434; spiniforme, 157, 434

Rhizoids, septate in Hepaticae, 57

Rhodobryum beyrichianum, 157; minutulum, 6

Rhynchostegiella compacta, 7

Rhynchostegiopsis auricolor, 164; flexuosa, 164, 398

Rhynchostegium sp., soluble sugars in, 471*

Rhynchostegium aquaticum, 165; brachypyxis, 165; debile, 440; nigrescens, 440; obscurum, 440; pulchellum, 165; scariosum, 166; semiscabrum, 166; serrulatum, 166; vitianum, 440

Riccardia latifrons, 8, 596; multifida, 596; palmata, 8; pinguis, 7, 281-; sinuata, 8, 596

Riccia albida, 597; bifurca, 597; californica, 281-; campbelliana, 281-; cilifera, 281-; crustata, 281-; crystallina, 224, 226*, 227*; crystallina, distribution in south Australia, 225*; curtisii, 597; donnellii, 597; fluitans, 50, 597; glauca, 50; gougetiana, 281-; hirta, 597; membranacea, 597; sorocarpa, 281-, 597; sullivantii, 597; trichocarpa, 281-

Ricciocarpus natans, 50, 597

Riella affinis, 281-; americana, 281-

Rigodium gracile, 162

Rinodina sect. Eurinodina, 129; sect. Placothallia, *130*

Rinodina, 82-, 181; balanina, 128-; californica, 131; diffractella, *139*; exigua, 590; hallii, 590; hueana, *132*; metaboliza, 516; mniaraea, 474, 590; mulleri, *139*; nimbosa, 128-; novomexicana, *132*; oreina, 128-; pachysperma, 567; physciaeformia, *139*; pyrina, 245, 414, 516; radiata, *135*; radiata subsp. lactea, *135*; stellata, *140*; suboreina, *132*; suboreina var. novomexicana, *132*; thysanota, *138*; turfacea, 590

Robinson, H. *See* Hermann, F. J., and H. Robinson

Rocky Mountains, lichen additions, 41

Roellia roellii, 109-

Ross Island, Antarctica, map of, 111*

Rundel, P. W.: Water relations and morphological variation in Ramalina menziesii, 23; (review), V. H. Heywood (editor), Taxonomy and Ecology. Proceedings of an International Symposium Held at the Department of Botany, University of Reading, 660; and P. A. Bowler: The lichen genus Trichoramalina, 188; *see* Bowler, P. A. and P. W. Rundel

Salazinic acid, 254, 637

Sarcogyne bicolor, 590; simplex, 590

Scapania, 628-; irrigua, 236-; lapponica, 236-; mucronata subsp. praetervisa, 74-; nemorosa, 281, 597; simmonsii, 195-

Schistidium apocarpum, 155

Schistochila appendiculata, SEM of spores and elaters, 299, 301*

Schistochila blumei, 57-
Schliephackea meteorioides, 153
Schlotheimia oerstediana, 159; rugifolia, 159; sublaevifolia, 159
Schuster, R. M. (review), H. Inoue, Illustrations of Japanese Hepaticae, 659
Scopelophila ligulata, 154
Seabury, F. *See* Magill, R. E., F. Seabury, and D. M. J. Mueller
Sekikaic acid, 617
Seligeria kunzeana, 152
Selkirk, D. R.: A fossil epiphyllous moss from the Miocene, 249
SEM methods: critical point drying method for liverwort spores, 284; evaluation of critical point drying technique in moss systematics, 628; fixation of liverwort gametophytes, 612; freeze-drying of liverwort spores, 284; preparation of liverwort spores and elaters, 283; preparation of soredia, 208
SEM studies of elaters in: Acrobolbus cinerascens, 299, 304*; Asterella tenera, 314, 316*, 318*; Blasia pusilla, 309*, 310; Chandonanthus squarrosus, 295, 298*; Cuspidatula monodon, 295, 300*; Frullania falciloba, 303, 306*, 307*; Haplomitrium gibbsiae, 287*; Herberta alpina, 288, 289*; Jungermannia hodgsoniae, 295, 296*; Lepicolea attenuata, 288, 290*, 292*; Lepidolaena clavigera, 291, 293*, 294*; Lepidozia spinosissima, 295, 297*; Marchantia berteroana, 317, 319*; Monoclea forsteri, 311*, 312; Neohodgsonia mirabilis, 320, 321*, 322*; Porella elegantula, 299, 305*; Schistochila appendiculata, 299, 301*; Targionia hypophylla, 313*, 314, 315*; Tetracymbaliella cymbalifera, 299, 302*; Treubia lacunosa, 308*, 310
SEM studies of leaves in: Desmatodon convolutus, 630*; Thelia asprella, 629*
SEM studies of spores in: Acrobolbus cinerascens, 299, 304*; Asterella tenera, 314, 316*, 318*; Blasia pusilla, 309*, 310; Chandonanthus squarrosus, 295, 298*; Cuspidatula monodon, 295, 300*; Fossombronia alaskana, 67*; Frullania falciloba, 303, 306*, 307*; Haplomitrium gibbsiae, 285*, 287*; Herberta alpina, 288, 289*; Jungermannia hodgsoniae, 295, 296*; Lepicolea attenuata, 288, 290*, 292*; Lepidolaena clavigera, 291, 293*, 294*; Lepidozia spinosissima, 295, 297*; Marchantia berteroana, 317, 319*; Monoclea forsteri, 311*, 312; Neohodgsonia mirabilis, 320, 321*, 322*; Porella elegantula, 299, 305*; Riccia crystallina, 227*; Schistochila appendiculata, 299, 301*; Targionia hypophylla, 313*, 314, 315*; Tetracymbaliella cymbalifera, 299, 302*; Treubia lacunosa, 308*, 310
SEM studies of water conducting elements in: Haplomitrium gibbsiae, 612, 613*, 615*; Haplomitrium rotundifolium, 615*; Hymenophyllum flabellatum, 612, 614*, 615*; Moerckia flotoviana, 615*; Symphyogyna circinata, 615*; Takakia ceratophylla, 615*; Takakia lepidozioides, 615*
Sematophyllum adnatum, 646; auratum, 167; caespitosum, 167, 468; cochleatum, 167; cuspidatum, 167; cuspidiferum, 167; **dimorphum** comb. nov., 167; entodontoides, 441; entodontoides var. **moorei** stat. nov., 441; flavidum, 167; galipense, 167; **harpidioides** comb. nov., 167; hawaiiense, 441; insularum, 167; kegelianum, 167; lindigii, 167; loxense, 167; moorei, *441*; pungens, 167; sericefolium, 167; **simulans** comb. nov., 168; **subcespitosum** comb. nov., 168; subscabrum, 168; subsimplex, 168; **tonduzii** comb. nov., 168; xylophilum, 167
Seppelt, R. D.: Riccia crystallina in south Australia, 224
Sheard, J. W.: The genus Dimelaena in North America north of Mexico, 128; and M. E. Jonescu: A multivariate analysis of the distribution of lichens on Populus tremuloides in west-central Canada, 514
Sheel, *see* Chopra, R. N. and Sheel
Sigal, L. L. and D. R. Toren: New distribution records of lichens in California, 469
Skorepa, A. *See* Reyes, C. R. and A. Skorepa
Smiley, D. and C. J. George: Photographic documentation of lichen decline in the Shawangunk Mountains of New York, 179
Snider, J. A. and F. D. Bowers: Archidium, a moss genus new to Mexico, 639
Solberg, Y. J.: Studies on the chemistry of lichens. XII. Chemical investigations of the lichen species Xanthoria parietina (L.) Th. Fr.—2, 203
Solmsiella biseriata, 593-
Solorina crocea, 590; saccata, 590
Soredia, a new type in the Heppiaceae, 208; formation of, 210
South America, *see* Latin America
Sphaerocarpos, 63-; cristatus, 281-; donnellii, 281-; drewei, 281-; michelii, 281-; texanus, 281, 597
Sphaerophorus fragilis, 590; globosus, 590
Sphaerotheciella, generic characters, 533*
Sphagnum, 470-; capillaceum, 8, 151; costaricense, 151; cuspidatum, 151; erythrocalyx, 151; imbricatum, 647; magellanicum, 151, 644; meridense, 151; platycladum, 151;

plumulosum, 151; recurvum, 151; recurvum var. tenue, 84; squarrosum, 7; subsecundum, 151; subsecundum var. rufescens, 647; teres, 84; tonduzii, 151
Spiridens armatus, 435; balfourianus, 435; balfourii, 435
Splachnaceae gametangia: Splachnobryum flaccidum, 623; geheebii, 623; indicum, 623; obtusum, 623, 624*; synoicum, 623; Splachnum sphaericum, 623, 626*; wormskioldii, 626*; Tetraplodon angustatus, 623; mnioides, 623; Tayloria froelichiana, 623; lingulata, 623; serrata, 623
Splachnobryum gametangia, *see* Splachnaceae
Splachnobryum indicum, 433; obtusum, 156
Splachnum gametangia, *see* Splachnaceae
Splachnum ampullaceum, 33-; vasculosum, 74-
Sporastatia testudinea, 590
Spores: and elaters of liverworts, 281; scanning electron microscopy of, *see* SEM; measurement, statistical method for, 130; types of in Jungermanniales, 282; viability of in Desmatodon randii, 476
Sporophytes of Buxbaumia, developmental stages of, 504
Sporopollenin in liverworts, 324
Squamidium crispipilum, 161; isocladum, 161; leucotrichum, 161; longipilum, 161; macrocarpum, 161; nigricans, 161
Stableria osculatiana, 156
Staurothele clopima, 590; glacialis, 591; perradiata, 591
STEERE, W. C.: The occurrence of Mnium spinosum in North America. II., 79; Grimmia (Coscinodon) arctolimnia, a new species from Great Bear Lake, Northwest Territories, Canada, 230; (review), Y. Vasari, H. Hyvärinen, and S. Hicks (editors), Climatic changes in Arctic areas During the Last Ten-thousand Years. A Symposium Held at Oulanka and Kevo, 4–10 October, 1971, 103; and G. R. BRASSARD: The systematic position and geographical distribution of Fissidens arcticus, 195; and H. INOUE: Fossombronia alaskana, a new hepatic from Arctic Alaska, 63; and B. M. MURRAY: The geographical distribution of Bryum wrightii in arctic and boreal North America, 172
Stegonia latifolia, 63-
Stenodictyon nitidum, 398, 399*; pallidum, 400; sericeum, 164, 399*, 400
Stereocaulon albicans, 591; alpinum, 591; condensatum, 591; depreaultii, 591; glareosum, 591; intermedium, 591; paschale, 591; rivulorum, 591; subalbicans, 591; tomentosum, 591; vesuvianum, 591
Stereocaulon paschale, soluble sugars in, 466
Stereodon hamatus, 168
Stereophyllum, 441; affine, 166; contorteoperculatum, 166; cultelliforme, 166; leucostegum, 166; radiculosum, 166, 646; subobtusum, 166
Sticta limbata, 212; weigelii, 591
Stictic acid, 128-
Stokesiella oregana, 166; praelonga, 166
Strangospora, 82-, 220
Streptopogon calymperes, 155; erythrodontus, 155; rigidus, 155
Strigula elegans, 248; elegans var. antillarum, 248; nemathora, 248; nemathora var. hypothelia, 248; nitidula, 248; schizospora, 248; subtilissima, 248
STRINGER, M. H. L. *See* STRINGER, P. W. and M. H. L. STRINGER
STRINGER, P. W. and M. H. L. STRINGER: Studies on the bryophytes of southern Manitoba. VI. An ecological study of the bryophytes of coniferous forests in Bird's Hill Provincial Park, 1; Seventeen lichens new to Manitoba, 243; Air pollution and the distribution of epiphytic lichens and bryophytes in Winnipeg, Manitoba, 405; A quantitative study of corticolous bryophytes in the vicinity of Winnipeg, Manitoba, 551
STROEMER, J. R. *See* MELSTROM, C. E., N. C. MARAVOLO, and J. R. STROEMER
Südamerika, *see* Latin America
Sugars, soluble: in lichens, 464; in mosses, 470, 471*
Symblepharis helicophylla, 153; oerstediana, 153; vaginata, 153
Symphyogyna circinata, 615*
Symphysodon vitianus, 437
Symphysodontella cylindracea, 437
Synecology, Dicranella riparia, 236
Syrrhopodon subg. Orthotheca, 242-
Syrrhopodon aculeo-ciliatus, 154; apertus, 432; aristifolius, 432; banksii, 432; bernoullii, 154; berteroanus, 154; ciliatus, 433; ciliolatus, *242*; cristatus, 154; erubescens, 154; fissipapillatus, 433; flavescens, 154; gaudichaudii, 154; incompletus, 154, 241, 242; involutus, 433; lycopodioides, 154; mammillatus, 433; **mooreae** comb. nov., 433; parasiticus, 154, 242; prolifer, 154; quintasii, 242; rigescens, 433; subulatus, 433; texanus, 647; theriotii, 154; tristichellus, 433
Syzygiella perfoliata, 644

Takakia, archegonia, 627
Takakia ceratophylla, 615*; lepidozioides, 615*

Tapellaria epiphylla, 248
Targionia hypophylla, SEM of spores and elaters, 313*, 314, 315*
Targionia hypophylla, 49, 281-
Taxiphyllum planissimum, 166
Taxithelium falcifolium, 441; laxiusculum, 168; mundulum, 441; planum, 168; tenuisetum, 441; trachaelocarpum, 441; vernieri, 441
TAYLOR, J., P. J. HOLLINGSWORTH, and W. C. BIGELOW: Scanning electron microscopy of liverwort spores and elaters, 281
Tayloria gametangia, *see* Splachnaceae
Tayloria moritziana, 156; scabriseta, 156; splachnoides, in Arizona, 636
TEDESCHI, S. *See* MEYER, M. W., J. GREENBERG, and S. TEDESCHI
Telaranea nematodes, 597
TEM, methods of liverwort fixation, 612; preparation of soredia, 208
TEM studies of water conducting elements in Haplomitrium gibbsiae and Hymenophyton flabellatum, 612, 613*, 614*, 615
Temperature acclimation of Fontinalis novaeangliae, 17, 19*, 20*
Temporal variation of Buxbaumia aphylla sporophyte populations, 508*
Tetracymbaliella cymbalifera, SEM of spores and elaters, 299, 302*
Tetraphis pellucida, 6
Tetraplodon gametangia, *see* Splachnaceae
Tetraplodon mnioides, 156, 172-
Texas mosses, 645
Thamniopsis pendula, 164, 401*, 402
Thamnium exigua, 438; tumidicaule, *162*
Thamnobryum **tumidicaule** comb. nov., 162
Thamnolia subuliformis, 591; vermicularis, 591
Thelia asprella, SEM and use of critical point drying technique, 629*; 646
Thelotrema lepadinum, 591
Tholurna dissimilis, 591
Thuidiopsis furfurosa, 165, 440
Thuidium antillarum, 165; costaricense, 165; delicatulum, 8, 165, 646; delicatulum var. radicans, 6; erectum, 165; furfurosa, 165; furfurosum, 440; involvens, 165; leskeaefolia, 165; minutulum, 6, 165; miradoricum, 165; miradoricum var. gracilescens, 165; pellucens, 165; philibertii, 165; pseudo-protense, 165; ramosissimum, 440; recognitum, 8, 165; recognitum subsp. philibertii, 165; schistocalyx, 165; tahitense, 440; tamariscinum var. mexicanum, 165; urceolatum, 165; wrightii, 165
Thyridium constrictum, 433; constrictum var. tahiticum, 433; obtusifolium, 433
Thysanomitrium blumii, 430; richardii, 153; umbellatus, 430
Timmia megapolitana var. megapolitana, 6
Tomenthypnum nitens, 8, 74
Toninia aromatica, 591; ruginosa, 474, 591; squalida, 591
TOREN, D. R. *See* SIGAL, L. L. and D. R. TOREN
Tortella guatemalensis, 155; humilis, 646
Tortula caroliniana, 155; denticulata, 644; lacerifolia, 644; mniifolia, 155; mucronifolia, 63-, 415, 555; rhizophylla, 644; ruralis, 74-
Tortula princeps, sugars in, 471*
Trachyloma indicum, 437; tahitense, 437
Trachypus bicolor var. viridulus, 160
Trapelia coarctata, 591
TRAVERSE, A. (review), S. Nilsson (editor), World Pollen and Spore Flora, 495
Trematodon brevicollis, 77; latinervis, 431; longicollis, 152, 431; puteensis, 431; rapaensis, 431
Treubia lacunosa, SEM of spores and elaters, 308*, 310
Treubia nana, archegonia, 627
Tricharia albostrigosa, 248; carnea, 248; vulgaris, 248
Trichoramalina gen. nov., 191; climate, 190*; distribution, 193*
Trichoramalina **crinata** comb. nov., 191*, 192*; **melanothrix** comb. nov., 192*
Trichosteleum fluviale, 168; hamatum, 441; microcarpum, 168; orthophyllum, 442; patens, 442; pygmaeum, 442; verrucosum, 168
Trichostomopsis australasiae, 645
Trichostomum brachydontium, 155; campylocarpum, 155; **clavinerve** comb. nov., 433; oblongifolium, 433; tenuirostre, 155
Trichothelium epiphyllum, 248
Tristichium mirabile, 151

Ulota curvifolia, 74-
Umbilicaria angulata, 591; cylindrica, 591; deusta, 591; havaasii, 591; hyperborea, 591; lambii, 591; mammulata, 184; phaea, 591; proboscidea, 591; torrefacta, 591; vellea, 591
Unions: Amblystegium juratzkanum, 8; Aulacomnium palustre-Sphagnum capillaceum, 8; Mnium cuspidatum, 8; Mnium rugicum-Thuidium recognitum, 8
Usneaceae, 234-
Usnea comosa, 572; florida, 591; glabrescens, 591; hirta, 571, 591; longissima, 591; plicata, 591; trichodea, 591
Usnic acid, 128-, 234, 253-, 617, 637

Venezuela, *see* Latin America
Verrucaria acrotella, 592; hydrela, 592; nigrescens, 592
Vesicularia, 440; amphibola, 168; aperta, 442; bryifolia, 442; calodictyon, 442; crassicaulis, 168; inflectens, 442; longifolia, 443; subdenticulata, 168; tahitensis, 443
Vetaforma, 57
Voitia hyperborea, 172-

Washington, lichens of the North Cascades Range, 582, 583*
Water conducting elements, SEM and TEM studies in Haplomitrium gibbsiae and Hymenophyton flabellatum, 612, 613*, 614*, 615*
Water content in lichens, 25*
WEBER, W. A. (review), S. Flowers, Mosses: Utah and the West, 276; *see* NASH, T. H. and W. A. WEBER
Webera papillosa, 156, spectabilis, 156
Weissia clavinervis, 433; controversa, 433, 646; senocarpa, 154; viridula, 433
WELCH, W. H.: The Hookeriaceae of Central America, 328
WETMORE, C. M.: New type of soredium in the lichen family Heppiaceae, 208; (review), W. Martin and J. Child, Lichens of New Zealand, 104
WHITTIER, B. A. *See* WHITTIER, H. O. and B. A. WHITTIER
WHITTIER, H. O. and B. A. WHITTIER: List of mosses of southeastern Polynesia, 427
Wiesnerella denudata, 50
Wijkia costaricensis, 167; subnitidum, 167
Wilsoniella jardini, 431

Xantholamine, 204, 205*, 206*, 207*
Xanthoria, 208-, 561-; candelaria, 414, 567, 592; elegans, 128-, 137*, 592; fallax, 82-, 212, 213*, 245, 414, 516, 567; parietina, chemistry, 203, 204*; polycarpa, 414, 516, 592
Xenochila integrifolia, 57-

Zygodon intermedius, 436; liebmanii, 158; pungens, 158; pusillus, 154; reinwardtii, 158, 436; reinwardtii var. subintegrifolius, 436

Through the laburnum's dropping gold
Rose the light shaft of Orient mould,
And Europe's violets, faintly sweet,
Purpled the mossbeds at its feet.

FELICIA DOROTHEA HEMANS
The Palm Tree